

# WILLOWOOD USA

# WILLOWOOD OXYFLO 2EC

**ACTIVE INGREDIENT:**

Oxyfluorfen: 2-chloro-1-(3-ethoxy-4-nitrophenoxy)

4-(trifluoromethyl)benzene..... 22.3%

**OTHER INGREDIENTS:** ..... 77.7%**TOTAL:** ..... 100.0%

Contains 2 pounds active ingredient per gallon. Contains petroleum distillates, xylene or xylene range aromatic solvents.

EPA Reg. No. 87290-8

EPA Est. No. 61842-CA-001

## Keep Out of Reach of Children **WARNING - AVISO**

If you do not understand the label, find someone to explain it to you in detail.  
(Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.)

### SHAKE WELL BEFORE USING

### FIRST AID

**If on skin or clothing:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
- Call a poison control center or doctor for treatment advice.

**If swallowed:**

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.

### NOTE TO PHYSICIAN

Contains petroleum distillates. Vomitting may cause aspiration pneumonia.

**Manufactured For:**

Willowood, LLC

385 Interlocken Crescent, Suite #240

Broomfield, CO 80021

**Net Contents:****2.5 Gallons**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**WARNING AVISO**

Causes skin irritation. Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Do not get on skin, in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

**Mixers, loaders and applicators using engineering controls (see Engineering Controls requirements below) must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (such as barrier laminate or viton) when mixing and loading
- Chemical-resistant apron when mixing and loading

**All other mixers, loaders, applicators and other handlers must wear:**

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (such as barrier laminate or viton)
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when exposed to the product concentrate

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:** Mixers and loaders supporting aerial applications to fallow land or ground applications to corn, cotton, or soybeans must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], and must:

- Wear the personal protective equipment required above for mixers/loaders using engineering controls
- Wear protective eyewear if the system operates under pressure, and
- Be provided and have immediately available for use in case of emergency, such as a broken package, spill, or equipment breakdown, coveralls and chemical-resistant footwear.

Handlers performing applications to corn must use an enclosed cab that meets the definition in the Worker Protection Standard for agricultural pesticides [40 CFR 170.240 (d) (5)] for dermal protection. In addition, such applicators must:

- Wear the personal protective equipment required above for applicators using engineering controls
- Be provided and must have immediately available for use in an emergency when they must exit the cab in the treated area: coveralls, chemical-resistant gloves, chemical-resistant footwear, and chemical-resistant headgear, if overhead exposure
- Take off any PPE that was worn in the treated area before reentering the cab, and
- Store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (6)].

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when disposing of equipment wash water.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours except for the following:

- Onions, garlic and horseradish: The REI is 48 hours
- Conifer seedlings: The REI is 3 days
- Conifer trees: The REI is 6 days

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries or greenhouses.

**Do not enter or allow others to enter until sprays have dried.**

## PRODUCT USE INFORMATION

Willowood OxyFlo 2EC is a selective herbicide for postemergence and preemergence residual weed control in labeled crops. Directions provided in the Product Use Information section of this label apply to all uses of this product. Use directions for listed crops are provided in the Crop-Specific Use Directions section of this label.

### Use Restrictions

**The following use restrictions apply to all labeled uses of Willowood OxyFlo 2EC (Refer to directions for use for individual crops for additional crop-specific use restrictions.):**

- Do not graze or harvest plants from areas treated with Willowood OxyFlo 2EC for feed or forage.
- Apply Willowood OxyFlo 2EC only with ground equipment unless otherwise specified in crop-specific use directions.
- Willowood OxyFlo 2EC is phytotoxic to plant foliage. Avoid accidental spray contact or drift with established crops. Do not apply when weather conditions favor drift to non-target areas.
- Some labeled crops are tolerant to over-the-top applications of Willowood OxyFlo 2EC if applied during dormancy. Do not make over-the-top applications unless specifically allowed in crop-specific use directions.
- Do not treat ditch banks or waterways with Willowood OxyFlo 2EC or contaminate water used for irrigation or domestic purposes.
- Do not apply Willowood OxyFlo 2EC in enclosed greenhouse as foliage injury will result.

### Spray Drift Buffer Restrictions

- A 25 foot vegetative buffer strip must be maintained between all areas treated with this product and lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not allow spray to drift from the application site and contact people, structures people may occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.
- For ground boom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy when wind speed is 10 mph or less at the application site as measured by an anemometer.
- Use coarse spray according to ASAE 572 definition for standard nozzles or VMD of 475 microns for spinning atomizer nozzles.
- The applicator also must use all other measures necessary to control drift.

### Rotation Crop Restrictions

- Do not rotate to small-grain crops (includes barley, buckwheat, corn, pearl millet, proso millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, wild rice) within 10 months following an application of Willowood OxyFlo 2EC.
- Do not direct seed any crop, other than a crop labeled for use with Willowood OxyFlo 2EC, within 60 days following application.
- Do not transplant seedlings of crops, other than crops labeled for use with Willowood OxyFlo 2EC, within 30 days following application.
- **IMPORTANT: Unless otherwise specified elsewhere in this label, supplemental label or product bulletin, treated soil must be thoroughly mixed to a depth of 4 inches after harvest (or abandoning) of the treated crop but prior to planting of the rotational crop. Failure to achieve thorough and complete mixing or to follow the required minimum plant-back interval may result in crop injury, stand reduction and/or vigor reduction of the plant-back crop.** See specific fallow bed labeling instructions for required treatment-to-planting intervals following application of Willowood OxyFlo 2EC to fallow beds or fallow fields.

## Weeds Controlled

<b>Common Name</b>	<b>Scientific Name</b>
ageratum	<i>Ageratum conyzoides</i>
amaranth, spiny	<i>Amaranthus spinosus</i>
balsamapple	<i>Momordica charantia</i>
barnyardgrass, (watergrass)*	<i>Echinochola crus-galli</i>
bedstraw, catchweed	<i>Echinocholoa crus-galli</i>
bittercress, lesser	<i>Cardamine oligosperma</i>
bluegrass, annual*	<i>Poa annua</i>
buckwheat, wild	<i>Polygonum convolvulus</i>
burclover	<i>Medicago hispida</i>
buttercup, smallflower	<i>Ranunculus abortivus</i>
buttonweed	<i>Borreria laevis</i>
camphorweed	<i>Heterotheca subaxillaris</i>
canarygrass (annual)	<i>Phalaris canariensis</i>
carpetweed	<i>Mollugo verticillata</i>
cheeseweed (malva)	<i>Malva parviflora</i>
clover, red*	<i>Trifolium pratense</i>
clover, white*	<i>Trifolium repens</i>
cocklebur, common	<i>Xanthium pensylvanicum</i>
crabgrass, large (hairy)*	<i>Digitaria sanguinalis</i>
crotalaria	<i>Crotalaria species</i>
croton, tropic	<i>Croton glandulosus</i>
cudweed, narrowleaf	<i>Gnaphalium falcatum</i>
eveningprimrose, cutleaf	<i>Oenothera laciniata</i>
fiddleneck, coast*	<i>Amsinckia intermedia</i>
filaree, broadleaf	<i>Erodium botrys</i>
filaree, redstem	<i>Erodium cicutarium</i>
filaree, whitestem	<i>Erodium moschatum</i>
fireweed (from seed)	<i>Epilobium angustifolium</i>
flixweed	<i>Descurainia sophia</i>
foxtail, giant*	<i>Setaria faberi</i>
foxtail, green	<i>Setaria viridis</i>
foxtail, yellow	<i>Setaria lutescens</i>
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass*	<i>Eleusine indica</i>
groundcherry, cutleaf	<i>Physalis angulata</i>
groundcherry, Wright	<i>Physalis wrightii</i>
groundsel, common	<i>Senecio vulgaris</i>
henbit	<i>Lamium amplexicaule</i>
horseweed (marestail)	<i>Conyza canadensis</i>
jimsonweed	<i>Datura stramonium</i>

<b>Common Name</b>	<b>Scientific Name</b>
johnsongrass, seedling	<i>Sorghum halepense</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
ladysthumb (smartweed)	<i>Polygonum persicaria</i>
lambsquarters, common	<i>Chenopodium album</i>
lettuce, prickly (china lettuce)	<i>Lactuca serriola</i>
mallow, little (malva)	<i>Malva parviflora</i>
mayweed (dog fennel)	<i>Anthemis cotula</i>
minerslettuce	<i>Montia perfoliata</i>
morningglory species, annual	<i>Ipomoea species</i>
morningglory, ivyleaf*	<i>Ipomoea hederacea</i>
morningglory, tall*	<i>Ipomoea purpurea</i>
mustard, black	<i>Brassica nigra</i>
mustard, blue (purple mustard)	<i>Chorispora tenella</i>
mustard, common yellow	<i>Brassica campestris</i>
mustard, hedge	<i>Sisymbrium officinale</i>
mustard, tumble (Jim hill mustard)	<i>Sisymbrium altissimum</i>
mustard, wild	<i>Brassica kaber</i>
nettle, burning	<i>Urtica urens</i>
nightshade, American black	<i>Solanum americanum</i>
nightshade, black	<i>Solanum nigrum</i>
nightshade, hairy	<i>Solanum sarrachoides</i>
oats, wild	<i>Avena fatua</i>
orach, red	<i>Atriplex rosea</i>
oxalis (bermuda buttercup)	<i>Oxalis pes-caprae</i>
panicum, fall	<i>Panicum dichotomiflorum</i>
pepperweed, Virginia	<i>Lepidium virginicum</i>
pepperwood, yellowflower	<i>Lepidium perfoliatum</i>
pigweed, prostrate	<i>Amaranthus blitoides</i>
pigweed, redroot	<i>Amaranthus retroflexus</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
poinsettia, wild	<i>Euphorbia heterophylla</i>
puncturevine	<i>Tribulus terrestris</i>
purslane, common	<i>Portulaca oleracea</i>
pusley, florida	<i>Richardia scabra</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>
redmaids	<i>Calandrinia caulescens</i>
rocket, London	<i>Sisymbrium irio</i>
ryegrass, Italian	<i>Lolium multiflorum</i>
sage, lanceleaf	<i>Salvia reflexa</i>
sandbur, field	<i>Cenchrus inertus</i>
sandspurry, red	<i>Spergularia ruba</i>

<b>Common Name</b>	<b>Scientific Name</b>
sesbania, hemp	<i>Sesbania exaltata</i>
shepherdspurse*	<i>Capsella bursa-pastoris</i>
sicklepod	<i>Cassia obtusifolia</i>
sida, prickly (teaweed)	<i>Sida spinosa</i>
signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
smartweed, pennsylvania	<i>Polygonum pennsylvanicum</i>
sorrel, red (from seed)	<i>Rumex acetosella</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
speedwell, birdseye	<i>Veronica persica</i>
spurge, garden	<i>Euphorbia hirta</i>
spurge, prostrate**	<i>Euphorbia supine</i>
spurge, spotted**	<i>Euphorbia maculate</i>
spurry, corn	<i>Spergula arvensis</i>
tansymustard	<i>Descurainia pinnata</i>
thistle, bull**	<i>Cirsium vulgare</i>
thistle, Russian	<i>Salsola kali</i>
velvetleaf	<i>Abutilon theophrasti</i>
witchgrass	<i>Panicum capillare</i>
witchweed	<i>Striga asiatica</i>
woodsorrel, common yellow**	<i>Oxalis stricta</i>

\* Highest rate and/or multiple applications may be required for acceptable control.

\*\*Preemergence control only.

## **Application Methods and Cultural Practices**

### **Preemergence Weed Control**

Apply the specified rate in a broadcast spray volume of 15 or more gallons of water per acre using calibrated spray equipment capable of uniform application to the soil surface. Seedling weeds are controlled as they come in contact with the soil-applied herbicide during emergence. Preemergence weed control is most effective when Willowood OxyFlo 2EC is applied to soil surfaces that are clean (free of crop or weed residues or clippings) and weed-free. Prior to application, weed or crop residues should be removed by thorough incorporation into the soil using tillage equipment or by blowing the area to be treated. At least 0.25 inch of irrigation or rainfall is required to activate Willowood OxyFlo 2EC and should occur within 3 to 4 weeks after application. For optimum results, Willowood OxyFlo 2EC should be applied to prepared beds or soil surfaces that will be left undisturbed during the time for which weed control is desired. Cultural practices that disturb or redistribute surface soil following treatment with Willowood OxyFlo 2EC such as cutting water furrows will reduce weed control effectiveness.

**Application Rates and Rate Ranges:** Where rate ranges are given, use the lower rate range on coarse texture soils with less than 1% organic matter and lighter weed infestations. Use higher rates in the rate range on medium to fine texture soils, soils containing greater than 1% organic matter, heavy weed infestations, or for extended residual preemergence weed control.

### **Postemergence Weed Control**

Apply the specified rate in a broadcast spray volume of 20 or more gallons of water per acre (a minimum 10 gallons if applying Willowood OxyFlo 2EC in tank mix with glyphosate). Because Willowood OxyFlo 2EC is a contact herbicide, complete and uniform coverage of weed foliage is essential for optimum postemergence control. Increase the spray volume to ensure complete and uniform coverage as weed height and density increases or in the presence of heavy trash (weed or crop residue). Postemergence applications of Willowood OxyFlo 2EC are most effective when made to weeds at the seedling stage. Applications made later than the 4-inch or 4 leaf stage may result in partial control or suppression.

Postemergence applications should be made to seedling grasses not exceeding the 2-leaf stage. The addition of 0.25% v/v (2 pints per 100 gallons of spray) of an 80% active nonionic surfactant, labeled for application to growing crops, will enhance herbicidal effectiveness in controlling emerged weeds.

**Postemergence Application Rates:** Where a rate range is given, use a higher rate in the rate range for heavy weed infestations, weeds in advanced stages of growth or for extended residual preemergence weed control following control of existing emerged weeds.

**Ground Application**

**Ground Broadcast:** Apply Willowood OxyFlo 2EC using conventional low-pressure ground spray equipment with flat fan spray nozzles. Follow manufacturer’s recommendation for spraying pressure and boom height. An off-center (OC) nozzle positioned at the end of the boom may be desired. Check calibration of spray equipment before each use.

**Directed Sprays:** Apply Willowood OxyFlo 2EC as a coarse low-pressure spray in a spray volume of 20 or more gallons of spray per acre (broadcast basis). Follow manufacturer’s recommendations for nozzle spacing and operating pressure. Spray should be directed toward the soil at the base of the crop. In row crops, use a minimum of 2 flat fan nozzles per row (one on each side) and for optimum spray coverage use 4 flat fan nozzles per row (two on each side). The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and downward. With either sprayer system, nozzles should be adjusted to cover the weed foliage but minimize contact with the crop. Do not apply with hollow cone nozzles.

**IMPORTANT: Willowood OxyFlo 2EC is a contact herbicide. Contact of sprays or drift with foliage or green stems can cause severe crop injury. Use directed sprays and spray shields and/or leaf lifters as necessary to minimize contact of spray or drift with crop foliage or stems. Young green stems of woody plants are also susceptible to injury from spray contact. Potential for injury to woody stems diminishes with loss of green color and the development of relatively impervious non-living corky tissue (bark) on the surface of the stem.**

**Band Application:** Application rates listed in this label are for broadcast application. For band application, the rate per broadcast acre should be reduced according to the following formula:

$$\frac{\text{Band Width (in inches)}}{\text{Row Width (in inches)}} \times \text{Rate per Broadcast Acre} = \text{Amount Needed per Acre for Banded Application}$$

**Spot Application:**

For spot application, apply sprays uniformly to soil for preemergence weed control or on a spray-to-wet basis for postemergence weed control. Mix the required amount of Willowood OxyFlo 2EC with the recommended amount of water. For preemergence weed control, use one-half to one gallon of spray per 1000 sq ft. For postemergence weed control use a minimum of 1 gallon of spray per 1000 sq ft and add an 80% nonionic surfactant at the rate of 0.5 fl oz (1 Tbs) per gallon of spray. If making spot applications within an established crop, use coarse low-pressure sprays and direct the spray to the soil beneath the plants. To avoid crop injury, do not allow spray to contact leaves and stems of herbaceous plants or leaves or green stems of woody plants.

Amount of Willowood OxyFlo 2EC Required to Treat 1000 sq ft at Specified Application Rate					
0.5 pt/acre	1.0 pt/acre	2.0 pt/acre	3.0 pt/acre	4.0 pt/acre	8.0 pt/acre
0.2 fl oz. (5.5 ml)	0.4 fl oz. (11 ml)	0.75 fl oz. (22 ml)	1.1 fl oz. (33 ml)	1.5 fl oz. (44 ml)	3.0 fl oz. (88 ml)

1 pint = 16 fl oz.; 1 fl oz. = 29.6 (30) ml

**Aerial Application**

Use aerial boom equipment designed for use with herbicides and a minimum spray volume of 10 gallons per acre (5 gallons per acre if tank mixed with glyphosate). **Do not aerially apply Willowood OxyFlo 2EC unless crop-specific use directions specifically allow and provide directions for aerial application.**

**AVOID DRIFT: Exercise extreme care to avoid herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result. Extreme care must be exercised to prevent spray drift that could result in damage to other crops or desirable vegetation. Adhere to the following guidelines when aerial applications are to be made.**

**Spray Drift Management (Aerial Application):** Avoiding spray drift at the application site is the responsibility of the applicator. The potential for spray drift is controlled by the interaction of many equipment-and-weather-related factors. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry



formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator must adhere to the following requirements when Willowood OxyFlo 2EC is aerially applied:

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least 1/2 mile from all crops and desirable vegetation, except the following:

Maintain a minimum of downwind buffer zone of:

- 150 feet from dormant treefruit/nut/vine crops and overwintering sugar beets.
  - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets, and non-targeted vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
  4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any non-targeted vegetable fallow bed, crop, or desirable vegetation.

The use of a drift control agent may be required by local regulations. However, the drift control agent may decrease the weed control effectiveness.

**Important:** Aerial applicators must be familiar with the label for Willowood OxyFlo 2EC and follow all applicable use precautions. Applying Willowood OxyFlo 2EC in a manner other than recommended in this label is done at the user's risk. Users are responsible for all loss or damage resulting from aerial spraying. In addition, aerial applicators should follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, the most restrictive limitations apply.

### **For Aerial Application in Fresno County, California Only**

#### **(From February 15 through March 31 Only)**

In addition to the directions for use for aerial application appearing above, the following guidelines are required between the dates of February 15 and March 31 for applications in the following geographic area:

North: Fresno County line

South: Fresno County line

East: State Highway 99

West: Fresno County line

Observe the following directions to minimize off-site movement during aerial application of Willowood OxyFlo 2EC. Minimization of off-site movement is the responsibility of the grower, pest control advisor and aerial applicator.

A written recommendation must be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation must state the proximity of surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial application of Willowood OxyFlo 2EC is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to ensure that proper rates are being applied during the commercial use season. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation; or other written records showing calculations and measurements of flight and spray parameters acceptable.

Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

## Chemigation Instructions

Do not apply this product through any irrigation system unless the instructions for chemigation are followed. **Do not apply Willowood OxyFlo 2EC through chemigation equipment unless chemigation is allowed by Crop-Specific Use Directions.**

Apply this product only through sprinkler (center pivot, solid set, portable lateral, or low-volume (micro sprinkler)), drip (trickle), or flood (basin) irrigation systems. Refer to use directions for specific crops for instructions as to which type of irrigation system may be used. Do not apply this product through any other type of irrigation system.

- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## Sprinkler Chemigation (Foliar Spray Uses)

For sprinkler irrigation, sufficient water should be applied at the beginning of the irrigation period to ensure uniform wetting of the plant and/or soil surfaces. Meter Willowood OxyFlo 2EC into the sprinkler irrigation system at a continuous uniform rate during the middle 1/3 of the irrigation period to allow for uniform distribution to target weeds and/or soil surface. Continue irrigation during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. During sprinkler irrigation, sufficient water should be applied to ensure water penetration to a depth of two inches.

**AVOID DRIFT: Extreme care must be exercised to prevent spray drift that could result in damage to other crops or desirable vegetation. Use the following guidelines when applications of Willowood OxyFlo 2EC are made through sprinkler irrigation equipment.**

1. Do not apply when the wind direction is not stable, when inversion conditions exist, or when wind velocity exceeds 10 mph.
2. When wind speeds are 5 mph or less, maintain a minimum downwind buffer zone of at least 1/2 mile from all crops and desirable vegetation, except for the following:

Maintain a minimum downwind buffer zone of:

- 150 feet from dormant treefruit, dormant vines and overwintering sugar beets.
  - 650 feet from garlic, jojoba, legumes, onions, pastures, small grains, seedling sugar beets and vegetable fallow beds.
3. When wind speeds are between 5 and 10 mph, downwind buffer zones in excess of those listed above are suggested.
  4. For upwind and side borders, maintain a minimum buffer zone of 150 feet from any vegetable fallow bed, crop, or desired vegetation.

To apply pesticide using a sprinkler chemigation, the chemigation system must meet the following specifications.

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

#### **Flood (Basin) Chemigation (Soil Drench Uses)**

Willowood OxyFlo 2EC should be continuously metered into the water during the entire irrigation period. Agitation in the pesticide supply tank is suggested. Best weed control results from Willowood OxyFlo 2EC applied through flood (basin) irrigation systems are obtained when a uniform distribution and flow of irrigation water is maintained over level land.

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

#### **Drip (Trickle) Chemigation (Soil Drench Uses)**

To achieve optimum distribution of Willowood OxyFlo 2EC in the soil surface, meter Willowood OxyFlo 2EC at a continuous uniform rate during the middle 1/3 of the irrigation period. For best results, Willowood OxyFlo 2EC should be uniformly distributed across the wetted area to help reduce the “ring effect” of weed escapes. Continue irrigation during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system.

To apply a pesticide using drip (trickle) chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pipe and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when

the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

### Chemigation Calibration: For Low-Volume Sprinklers (Microsprinklers) and Drip (Trickle) Irrigation Systems

Calculation of use rate is based on wetted area around emitters- NOT on grove acres. To determine correct amount of Willowood OxyFlo 2EC, use the following formula:

1. Treated area per each emitter = A

$$A = 3.14 \times (\text{radius} \times \text{radius})$$

**Example:** If the average distance from emitter to perimeter of wetted area measured at the soil surface is 13 inches, then

$$A = 3.14 \times (13'' \times 13'')$$

$$A = 3.14 \times (169'')$$

$$A = 530.7 \text{ square inches}$$

2. The area in square feet wet in each acre = B

$$B = \frac{A \times \text{emitters/acre}}{144}$$

**Example:** If there are 300 emitters per acre, then

$$B = \frac{530.7 \times 300}{144} = B = 1105.6 \text{ square feet wetted per acre}$$

3. The total area (in square feet) wet by your system = C

$$C = B \times \text{acres covered by system}$$

**Example:** If the system covers 20 acres, then

$$C = 1105.6 \text{ square feet per acre} \times 20 \text{ acres}$$

$$C = 22,112 \text{ square feet wetted by system}$$

4. Amount of Willowood OxyFlo 2EC to inject = S

$$\text{Rate per treated acre of Willowood OxyFlo 2EC} = R$$

$$S = \frac{C \times R}{43,560} = \text{quarts of Willowood OxyFlo 2EC}$$

**Example:** If the desired application rate per treated acre is 1 quart of Willowood OxyFlo 2EC, then

$$S = \frac{22,112 \times 1.0}{43,560} = S = 0.507 \text{ quarts of Willowood OxyFlo 2EC should be injected into system.}$$

**Note:** Select the proper rate based on weed spectrum and desired length of control (See **Rate Ranges** section below).

### Chemigation Systems Connected to Public Water Systems

If the chemigation system is connected to a public water supply, the following conditions must also be met.

- Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the

supply tank when the irrigation system is either automatically or manually shutdown.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

### **Mixing Directions**

**Shake well before use.** Fill the spray tank at least one-third full of clean water. With the pump and agitator running, add the recommended amount of herbicides to the spray tank. The order of addition to the spray tank should be (1) wettable powders, (2) flowables and (3) soluble liquids. Complete filling of the spray tank with water.

**Use of Surfactants:** For all applications of Willowood OxyFlo 2EC where postemergence weed control is desired (except garlic and onions), add a minimum of 2 pints of 80% active nonionic surfactant (cleared for application to growing crops) per each 100 gallons of spray. The addition of 4 pints of nonionic surfactant is recommended to enhance postemergence activity when hard water (greater than 600 ppm) is used. Maintain agitation until spraying is completed.

### **Tank Mixing Precautions:**

- Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.
- Do not exceed recommended application rates. Do not tank mix this product with another pesticide that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.

**Tank Mix Compatibility Testing:** A jar test is recommended prior to tank mixing to ensure compatibility of this product and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the

jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jells, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

**Sprayer Clean-up:** Thoroughly flush spray equipment (tank, pump, hoses and boom) with clean water before and after each use. Residues of Willowood OxyFlo 2EC remaining in spray equipment may damage other crops. The addition of a non-ionic surfactant to equipment flushing waters at the rate of 1 quart per 100 gallons is recommended to aid in removal of residue of Willowood OxyFlo 2EC.

## CROP SPECIFIC USE DIRECTIONS

### ARTICHOKE (GLOBE)

#### Post-Directed Spray Application

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b>  <b>Postemergence</b>	4-6	<b>Application Method:</b> Apply as a directed spray to the soil surface between the rows and at the base of artichoke plants in a minimum spray volume of 40 gallons per acre.  <b>Timing to Crop:</b> Apply after completion of ditching operations. Separate applications of up to 4 pts/acre may be made 8 to 10 weeks apart or a single application of up to 6 pts/acre may be made.  <b>Timing to Weeds:</b> Preemergence up to 8 leaf stage.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Do not apply over-the-top. Contact with direct spray or drift will cause injury to artichoke fronds or severe injury to buds or flowers.</li> <li>• Application of Willowood OxyFlo 2EC to artichoke plantings should be delayed a minimum of 60 days after cutting back or transplanting.</li> </ul>		
<b>Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not apply more than 6 pints of Willowood OxyFlo 2EC per acre per season as a result of a single application or multiple applications.</li> <li>• <b>Preharvest Interval:</b> Do not apply within 5 days of harvest.</li> </ul>		

#### Key Weeds Controlled:

##### Preemergence

cheeseweed (malva)  
 groundsel, common  
 lambsquarters, common

##### Postemergence

cheeseweed (malva)  
 groundsel, common  
 mustard, common yellow

**Preemergence**

mustard, common yellow  
 oxalis (bermuda buttercup)\*  
 shepherdspurse  
 sowthistle, annual

**Postemergence**

nettle, burning  
 oxalis (bermuda buttercup)  
 shepherdspurse  
 sowthistle, annual

\*Suppression

**BROCCOLI/CABBAGE/CAULIFLOWER****Pre-transplant (Preplant) Application for Preemergence Broadleaf Weed Control**

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1-2	<p><b>Pre-Transplant Application Only:</b> Apply broadcast to final seedbed prior to transplanting. Use lower rate in the rate range on coarse textured soils with less than 1% organic matter. Use the highest rate range on medium to fine textured soils or soils containing greater than 1% organic matter.</p> <p>Transplanting should be accomplished with minimal soil disturbance and soil left undisturbed during the time weed control is desired.</p>

**Precautions:**

- Pre-transplant applications may result in initial, but temporary, crop injury (leaf cupping or crinkling) and is enhanced if crop leaves come in direct contact with treated soil. Crop will rapidly outgrow this condition and develop normally. Severe crop injury may result if transplants are under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides or storage conditions. The use of transplants less than 5 weeks old or use of extremely succulent transplants grown in containers less than 1 inch square, may increase the severity of crop injury. Hardening off, increasing the age of transplants or increasing the size of the rooting containers will lessen the possibility and/or severity of potential crop injury.
- Willowood OxyFlo 2EC will assist in early season annual grass control, however, a herbicide program for preemergence or postemergence control of annual grasses is recommended.
- Note: Do not apply Willowood OxyFlo 2EC if an acetanilide herbicide such as Dual Magnum herbicide, Lasso herbicide, or Ramrod herbicide has been applied to the field during the current growing season as severe crop injury may occur.
- Do not apply Willowood OxyFlo 2EC as a preemergence treatment to direct-seeded broccoli, cabbage or cauliflower.
- Do not apply Willowood OxyFlo 2EC post-transplant or over-the-top of broccoli, cabbage or cauliflower.
- Applications to muck soils may result in partial weed control or suppression.
- Furrow and drip irrigation immediately after transplanting and under high temperatures can result in increased crop injury. Sprinkler irrigation is recommended during early establishment of transplants. If these conditions cannot be met, Willowood OxyFlo 2EC herbicide should not be used.

**Crop-Specific Restrictions:**

- Do not apply more than 2 pints of Willowood OxyFlo 2EC per treated acre per season.

**Key Weeds Controlled:**

**Preemergence**

carpetweed

pigweed, redroot

purslane, common

smartweed, Pennsylvania

**CACAO (BEARING AND NONBEARING)**

**(For Use Only in Hawaii)**

Willowood OxyFlo 2EC may be applied as a pre-transplant treatment or to established or recently transplanted cacao.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b> <b>Postemergence</b>	2-8	<b>Pre-transplant Application:</b> Up to 4 pints per broadcast acre may be applied as a pre-transplant application. <b>Application to Established Plantings:</b> In established plantings, including recently transplanted cacao plants, apply as a directed spray to the orchard floor. Use higher rates in rate range and increase spray volume to control dense growth of existing weeds or for extended residual preemergence weed control.

**Precautions:**

- Do not apply preplant or preemergence to direct-seeded cacao.
- Willowood OxyFlo 2EC should be applied to only healthy growing trees/transplants of suitable size to allow directed sprays. Avoid spray contact with foliage.



Weed Control	Rate (pt/acre)	Specific Use Directions
<p><b>Crop-Specific Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 8 pints of Willowood OxyFlo 2EC per acre as a single application or more than 24 pints per acre per year</li> <li><b>Preharvest Interval:</b> Do not apply Willowood OxyFlo 2EC within 1 day of harvest.</li> </ul>		

**Key Weeds Controlled:**

**Preemergence**

ageratum  
 buttonweed  
 crotalaria  
 purslane, common  
 spurge, garden

**Postemergence**

purslane, common  
 spurge, garden

**CITRUS (NONBEARING)**

**Citrus, such as Calamondin, Chironja, Citrus Citron, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Pummelo, Satsuma Mandarin, Sour Orange, Sweet Orange, Tangelo, Tangerine, Tangor**

Willowood OxyFlo 2EC may be applied only in non-bearing citrus orchards. Apply only as a directed spray to the orchard floor avoiding contact with citrus foliage.

Weed Control	Rate (pt/acre)	Specific Use Directions
--------------	----------------	-------------------------

Specimen

<b>Preemergence</b>	6	<p><b>Preemergence Weed Control:</b> Up to 6 pts/acre may be applied for residual preemergence weed control.</p> <p><b>Postemergence Weed Control:</b> The 6 pint/acre rate will control weeds up to 4 inches tall. Weeds greater than 4-leaf or 4 inches tall may be partially controlled. Use sufficient spray volume for complete and uniform coverage of weeds. Increase the spray volume with increased weed height and density to ensure complete coverage.</p>
<b>Postemergence</b>	2-6	
<p><b>Tank Mixing:</b> Refer to Mixing Directions section for Tank Mixing Precautions.</p> <ul style="list-style-type: none"> <li>• <b>Preemergence Use:</b> For residual control of grass weeds, Willowood OxyFlo 2EC may be tank mixed with grass herbicides labeled for use in citrus.</li> <li>• <b>Postemergence Use:</b> For broader spectrum postemergence control of emerged grass and broadleaf weeds, Willowood OxyFlo 2EC may be tank mixed with paraquat (Gramoxone herbicide) or glyphosate.</li> </ul>		
<p><b>Precautions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply during periods of new citrus foliage growth. Applications should be made after foliage has fully expanded and hardened off. Avoid direct spray contact with citrus foliage.</li> </ul>		
<p><b>Crop-Specific Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Apply Willowood OxyFlo 2EC only to nonbearing citrus (trees that will not bear fruit for one year).</li> <li>• Do not apply more than 6 pints of Willowood OxyFlo 2EC per acre per year as a result of single or multiple applications.</li> </ul>		

**Key Weeds Controlled:**

**(Arizona and California)****(Florida, Louisiana and Texas)****Preemergence**

burclover  
cheeseweed (malva)  
fiddleneck, coast  
filaree, broadleaf  
filaree, redstem  
filaree, whitestem  
groundsel, common  
henbit  
knotweed, prostrate  
lambsquarters, common  
lettuce, prickly  
pigweed, redroot  
purslane, common  
redmaids  
rocket, London  
shepherdspurse  
sowthistle, annual  
spurge, prostrate  
spurge, spotted

**Postemergence**

cheeseweed (malva)  
fiddleneck, coast  
filaree, broadleaf\*  
filaree, redstem\*  
filaree, whitestem\*  
groundsel, common  
henbit  
minerslettuce  
nettle, burning  
pigweed, redroot  
redmaids  
shepherdspurse  
sowthistle, annual

**Preemergence**

cudweed, narrowleaf  
eveningprimrose, cutleaf\*\*  
groundcherry, cutleaf  
lambsquarters, common  
nightshade, American  
black  
nightshade, black  
pepperwood, Virginia  
pigweed, redroot  
poinsettia, wild  
pusley, Florida  
sida, prickly (teaweed)  
smartweed, Pennsylvania  
sowthistle, annual  
spurge, prostrate  
spurge, spotted

**Postemergence**

balsamapple  
cudweed, narrowleaf\*\*\*  
eveningprimrose, cutleaf\*\*  
groundcherry, cutleaf  
groundcherry, Wright  
lambsquarters, common  
morningglory, annual  
nightshade, American  
black  
nightshade, black  
pepperweed, Virginia  
pigweed, redroot  
poinsettia, wild  
purslane, common  
pusley, Florida  
sida, prickly (teaweed)  
smartweed, Pennsylvania  
sowthistle, annual

\* Willowood OxyFlo 2EC at the 6 pt/acre will provide control of filaree and other weeds up to 4-inch stage. Applications to weeds beyond the 4-inch stage may result in partial control.

\*\*Highest rate and/or multiple applications may be required for acceptable control.

\*\*\*Maximum 0.5-inch diameter.

**CLARY SAGE****Clary Sage (*Salvia sclarea*) Grown and Utilized in the Essence Industry****(For Use Only in North Carolina)**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Postemergence</b>	0.5 – 1	Willowood OxyFlo 2EC may be applied to established clary sage for control of henbit ( <i>Lamium amplexicaule</i> ) and other winter annual broadleaf weeds during the winter and spring season.  Apply shortly after the first flush of henbit is in the 2- to 4-leaf stage of growth. Additional applications may be required to control subsequent weed flushes through the spring season. After treatment, henbit will stop growing and slowly die. Increase the spray volume if weed growth is dense.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Clary sage may respond to the topical application of this product with some marginal leaf burn, but recovery is rapid.</li> </ul>		
<b>Crop-Specific Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 6 pints per acre per year.</li> </ul>		

**COFFEE (BEARING AND NONBEARING)****(For Use Only in Hawaii)**

Willowood OxyFlo 2EC may be applied to established coffee, recently transplanted coffee, or as a pre-transplant treatment. In established non-dormant coffee, apply as a directed spray avoiding contact with crop foliage. Newly established transplants should be healthy and well established and of sufficient size to allow use of directed sprays without contacting crop foliage.

Willowood OxyFlo 2EC may be applied over-the-top of dormant coffee transplants. Transplants are considered to be dormant when active terminal growth has ceased and terminal buds have formed. Application over-the-top of coffee plants after buds start to swell (a sign that new growth has resumed) may result in crop injury and is not recommended.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b> <b>Postemergence</b>	2-8	<b>Preemergence Weed Control:</b> <ul style="list-style-type: none"> <li>• Apply as a directed spray to the orchard floor beneath established coffee plants.</li> <li>• Up to 4 pints per acre may be applied as a pre-treatment application prior to transplanting coffee plants.</li> </ul> <b>Postemergence Weed Control:</b> Increase the spray volume when weed growth is dense or trash is present; or use a higher rate within the rate range for extended residual preemergence weed control.
<b>Tank Mixing:</b> Refer to Mixing Directions section for Tank Mixing Precautions: Apply tank mixes only as a directed spray.		
<b>Precaution:</b> <ul style="list-style-type: none"> <li>• To prevent foliar injury, do not apply during periods of rapid new growth or allow spray to contact actively growing foliage.</li> </ul>		
<b>Crop-Specific Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not apply preplant or preemergence to direct-seeded coffee.</li> <li>• Do not apply more than 8 pints per broadcast acre of Willowood OxyFlo 2EC in a single application or 24 pints per broadcast acre per year.</li> <li>• <b>Preharvest Interval:</b> Do not apply Willowood OxyFlo 2EC within one (1) day of harvest.</li> </ul>		

**Key Weeds Controlled:**

**Preemergence**

ageratum  
 buttonweed  
 crotalaria  
 purslane, common  
 spurge, garden

**Postemergence**

purslane, common  
 spurge, garden

**CONIFER SEEDBEDS, TRANSPLANTS, CONTAINER STOCK AND SELECTED FIELD GROWN DECIDUOUS TREES**

**General Use Precautions and Restrictions:**

- Not for conifer release in forest management programs or for forest regeneration applications.
- Do not apply Willowood OxyFlo 2EC in an enclosed greenhouse structure as injury to plant foliage may result.
- Do not store or transport treated container stock in an enclosed structure until completion of 4 irrigations (minimum 21 days) as injury to non-labeled plants may occur.
- Apply Willowood OxyFlo 2EC only to healthy conifer stock. Do not apply Willowood OxyFlo 2EC to conifers that are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result.
- Do not graze or harvest livestock forage from treated areas.

**Key Weeds Controlled:** When Willowood OxyFlo 2EC is applied preemergence or postemergence at recommended

dosages and weed stages.

barnyardgrass*	mustard, blue
bedstraw, catchweed	mustard, tumble
bittercress, lesser	mustard, wild
bluegrass, annual*	nettle, burning
buckwheat, wild	nightshade, black
burclover	nightshade, hairy
carpetweed	oats, wild
clover, red*	orach, red
clover, white*	pepperweed, yellowflower
cocklebur, common	pigweed, prostrate
crabgrass, large*	pigweed, redroot
fiddleneck, coast*	pimpernel, scarlet
filaree, broadleaf	purslane, common
filaree, redstem	redmaids
fireweed (from seed)	rocket, London
flixweed	sandspurry, red
foxtail, giant*	shepherdspurse*
goosegrass*	sida, prickly
groundcherry, cutleaf	smartweed, Pennsylvania
groundcherry, Wright	sorrel, red (from seed)
groundsel, common	sowthistle, annual
henbit	speedwell, birdseye
jimsonweed	spurge, prostrate**
knotweed, prostrate	spurge, spotted**
ladysthumb	spurry, corn
lambsquarters, common	tansymustard
lettuce, prickly	thistle, bull**
mallow, little	thistle, Russian
mayweed	velvetleaf
minerslettuce	witchgrass
morningglory, ivyleaf*	woodsorrel, yellow**

morningglory, tall\*

\*Highest rate and/or multiple applications may be required for acceptable control.

\*\*Preemergence control only.

## CONIFER SEEDBEDS

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 3 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Willowood OxyFlo 2EC provides both postemergence and residual preemergence control of many broadleaf weeds and annual grass species.

Seeded conifers are tolerant to preemergence and postemergence applications of Willowood OxyFlo 2EC. For weed control during the establishment of conifer seedlings, Willowood OxyFlo 2EC can be applied after seeding of conifers, but prior to emergence. For weed control in emerged conifers, Willowood OxyFlo 2EC may be applied over-the-top, but application should be delayed a minimum of 5 weeks after seedling emergence. If application is made during cool, cloudy weather, make certain that seedlings have hardened-off prior to spraying.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	1-4	<b>Application after planting, but prior to emergence of conifer seedlings:</b> Where grass weeds are present, apply 2 to 4 pints of Willowood OxyFlo 2EC per acre. In known areas of high weed competition, apply 4 pints of Willowood OxyFlo 2EC per acre. Broadcast to beds and irrigate with 1/2 to 3/4 inch of sprinkler irrigation before weed emergence. Willowood OxyFlo 2EC is most effective on annual grasses when applied preemergence.
Postemergence	1-2	<b>Application after emergence of conifer seedlings:</b> Application should be made to seedling weeds less than 4 inches in height (seedling grasses not exceeding the 2-leaf stage). Depending on subsequent weed flushes, multiple applications may be necessary to achieve season-long weed control.

**Chemigation:** Willowood OxyFlo 2EC may be applied at labeled rates through sprinkler irrigation systems. For center pivot irrigation systems, apply the specified dosage of Willowood OxyFlo 2EC per acre metered at a continuous uniform rate during the entire irrigation period, otherwise meter Willowood OxyFlo 2EC at a continuous uniform rate during the middle 1/3 of the irrigation period. When applying by sprinkler irrigation, follow directions given in the Chemigation Instructions section of this label.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Occasionally spotting, crinkling, or flecking may appear on leaves of conifers. Leaves that receive direct spray or drift may be injured, but typically outgrow this condition rapidly and develop normally.</li> </ul>		
<b>Crop-Specific Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 8 pints of Willowood OxyFlo 2EC per acre per year.</li> </ul>		

**Willowood OxyFlo 2EC may be applied to conifer seedbeds of the following species:**

**Important:** When applied as directed, the conifer species listed on this label have shown tolerance to Willowood OxyFlo 2EC. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of listed species under all possible growing conditions. Until familiar with results under local growing conditions, the user should exercise reasonable judgment and caution with this product. Limit application of this product to a few plants in a small area to determine plant tolerance and extent of injury if such occurs, prior to initiating large-scale applications.

<b>Douglas Fir</b>	<i>Pseudotsuga menziesii</i>
<b>Fir</b>	Fraser ( <i>Abies fraseri</i> ) Grand ( <i>Abies grandis</i> ) Noble ( <i>Abies procera</i> )
<b>Hemlock</b>	Eastern hemlock ( <i>Tsuga canadensis</i> )
<b>Pine</b>	Austrian ( <i>Pinus nigra</i> ) Eastern White ( <i>Pinus strobes</i> ) Himalayan ( <i>Pinus wallichiana</i> ) Jack ( <i>Pinus banksiana</i> ) Loblolly ( <i>Pinus taeda</i> ) Lodgepole ( <i>Pinus contorta</i> ) Longleaf ( <i>Pinus palustris</i> ) Monteray ( <i>Pinus radiata</i> ) Mugho ( <i>Pinus mugo</i> ) Ponderosa ( <i>Pinus ponderosa</i> ) Scotch ( <i>Pinus sylvestris</i> ) Shortleaf ( <i>Pinus echinata</i> ) Slash ( <i>Pinus elliottii</i> ) Virginia ( <i>Pinus virginiana</i> )
<b>Spruce</b>	Blue ( <i>Picea pungens</i> ) Dwarf ( <i>Picea glauca Conica</i> ) Alberta ( <i>Picea abies</i> ) Norway ( <i>Picea sitchensis</i> )

**CONIFER TRANSPLANTS AND CONTAINER STOCK  
(INCLUDES 2-0 SEEDLING AND CHRISTMAS TREE PLANTINGS)**

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 6 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Many container-grown conifers and conifer transplants are tolerant to preemergence and postemergence applications of Willowood OxyFlo 2EC. Applied postemergence, Willowood OxyFlo 2EC provides postemergence control of emerged weeds and preemergence residual control of many broadleaf weed and grasses (See Key Weeds Controlled) at the beginning of this section.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b> <b>Postemergence</b>	4-8	<b>Transplanted and Container Grown Conifers:</b> For best results, preemergence applications should be made immediately after transplanting seedlings or to weed-free container stock. Postemergence applications should be made to weeds less than 4 inches in height. Two applications may be necessary, in fall-transplanted conifer fields, for season-long weed control. The addition of a non-ionic surfactant (0.25% v/v) labeled for application to growing crops, enhances the activity of Willowood OxyFlo 2EC on emerged weeds.
<b>Precautions:</b> <ul style="list-style-type: none"> <li>• Do not make over-the-top applications during periods of active conifer growth. Apply only before bud break or after new terminal growth has hardened off.</li> </ul>		
<b>Crop-Specific Restrictions:</b> <ul style="list-style-type: none"> <li>• Do not apply more than 8 pints of Willowood OxyFlo 2EC per acre in a single application or more than 16 pints per acre per year.</li> </ul>		

**In addition to those conifer species listed under the Conifer Seedbed section, the following conifer species have been shown to be tolerant to Willowood OxyFlo 2EC:**

- |                        |  |
|------------------------|--|
| <b>Arborvitae</b>      | <i>Thuja occidentalis</i><br><i>Thuia orientallis</i>  |
| <b>Juniper</b>         | <i>Juniperus chinensis</i><br><i>Juniperus horizontalis</i><br><i>Juniperus procumbens</i><br><i>Juniperus sabina</i><br><i>Juniperus scopulorum</i> |
| <b>Red Cedar</b>       | <i>Juniperus virginiana</i>  |
| <b>Western Hemlock</b> | <i>Tsuga heterophylla</i>  |
| <b>Yew</b>             | <i>Taxus species</i>   |

**SELECTED FIELD-GROWN DECIDUOUS TREES**

Listed field-grown deciduous trees are tolerant only to directed spray applications of Willowood OxyFlo 2EC. Willowood OxyFlo 2EC provides both preemergence and postemergence control of listed broadleaf weeds and grasses.

**Timing to Crop:** Apply Willowood OxyFlo 2EC to established deciduous trees or after transplanting. For optimum weed



control, applications should be made prior to weed germination. Apply only as a directed spray to soil beneath the trees.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence Postemergence</b>	2-6	<p>Willowood OxyFlo 2EC may be applied to established deciduous trees or after transplanting as a single or split application. Apply as a directed spray to the soil surface. Use of spray shields to reduce exposure of foliage and bark is recommended. The addition of a non-ionic surfactant (0.25% v/v) labeled for application to growing crops, will enhance herbicidal activity on emerged weeds.</p> <p><b>Spot Application:</b> Spot treatments at specified rates may be used to control localized weed infestations. See use directions for Spot Application in the Application Methods and Recommended Cultural Practices section.</p>
<p><b>Tank Mixing:</b> For broader spectrum control, Willowood OxyFlo 2EC may be tank mixed with other preemergence or postemergence herbicides registered for this use in deciduous trees. Refer to the Mixing Directions section for Tank Mixing Precautions.</p>		
<p><b>Precautions:</b></p> <ul style="list-style-type: none"> <li>• For maximum crop safety, directed applications should be prior to budbreak in the spring or after trees have initiated dormancy in the fall. Avoid contact of spray or drift with foliage or stems with green bark. Application after bud swell may result in crop injury and is not recommended. If a non-dormant application is required due to weed competition, apply only after foliage has fully expanded and hardened off. Use only directed sprays and spray shields to prevent spray contact with stems with green bark or foliage.</li> <li>• Do not apply Willowood OxyFlo 2EC to trees that have been weakened or are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, wind injury, drought, flooding, previously applied pesticides, insects, or water injury as severe injury may result.</li> </ul>		
<p><b>Crop-Specific Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 6 pints of Willowood OxyFlo 2EC per acre per year.</li> <li>• Do not apply to bearing treefruit, nut or vine crops. For selected bearing treefruit, nut and vine crops, refer to Treefruit/Nut/Vine section of this label for use directions.</li> <li>• Do not graze or feed livestock forage cut from areas treated with Willowood OxyFlo 2EC.</li> </ul>		

**Willowood OxyFlo 2EC may be applied to the following deciduous tree species:**

Almond**	<i>Prunus</i> spp.
Apple**	<i>Malus X domestica</i>
Apricot**	<i>Prunus</i> spp.
Ash, Green	<i>Fraxinus pennsylvanica</i>
Ash, White	<i>Fraxinus americana</i>

Birch, River

Cherry\*\*

Chestnut\*\*

Crabapple\*\*

Cottonwood

Dogwood

Eucalyptus

Filbert\*\*

Lilac

Locust, Black

Maple, Black\*

Maple, Red\*

Maple, Sugar\*

Myrtle, Crepe

Nectarine\*\*

Nut, Hickory\*\*

Nut, Macadamia

*Betula nigra*

*Prunus* spp.

*Castanea* spp.

*Malus* spp.

*Populus* spp.

*Cornus florida*

*Eucalyptus viminalis*

*Eucalyptus pulverulenta*

*Eucalyptus camaldulensis*

*Corylus* spp.

*Syringa vulgaris*

*Robinia pseudoacacia*

*Acer nigrum*

*Acer rubrum*

*Acer saccharum*

*Lagerstoemia indica*

*Prunus* spp.

*Carya* spp.

*Macadamia ternifolia*

Specimen

Oak, Chestnut	<i>Quercus prinus</i>
Oak, Cherrybark	<i>Quercus pagoda</i>
Oak, Nutt All	<i>Quercus nuttallii</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus rubra</i>
Oak, Water	<i>Quercus nigra</i>
Oak, Willow	<i>Quercus phellos</i>
Olive, Russian	<i>Elaeagnus angustifolia</i>
Poplar	<i>Populus</i> spp.
Poplar, Tulip	<i>Liriodendron tulipifera</i>
Peach**	<i>Prunus persica</i>
Pear**	<i>Pyrus</i> spp.
Pecan**	<i>Carya</i> spp.
Pistachio**	<i>Pistacia vera</i>
Plum**	<i>Prunus</i> spp.
Prune**	<i>Prunus</i> spp.
Redbud	<i>Cercis Canadensis</i>
Sweetgum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>
Walnut, Black**	<i>Juglans nigra</i>

\* Do not apply to maple trees used for production of maple sap or maple syrup.

\*\*Apply only to nonbearing trees. For bearing treefruit, nut and vine crops, refer to specific use directions in the Treefruit/ Nut/Vine section of this label.

**CORN**

**FOR USE ONLY ON FIELD CORN IN CONJUNCTION WITH THE USDA WITCHWEED ERADICATION PROGRAM IN NORTH CAROLINA AND SOUTH CAROLINA**

Apply Willowood OxyFlo 2EC only as a directed spray from May through August for preemergence and postemergence control of witchweed (*Striga asiatica*). Corn must be a minimum of 24 inches tall. Examine witchweed infested fields during the early part of the growing season to determine uniformity of corn stand and grass weed pressure. If necessary, cultivate weed-infested fields prior to initial application of Willowood OxyFlo 2EC to allow for optimum soil coverage during the initial application. Fields treated with Willowood OxyFlo 2EC should be inspected regularly for any breakthrough of witchweed. If breakthrough occurs, a second application should be made as soon as possible after appearance of witchweed. Repeat treatments should occur prior to bloom stage to prevent seed set.

Weed Control	Rate (pt/acre)	Specific Use Directions
--------------	----------------	-------------------------

<b>Preemergence</b>	2-3	<b>Initial Application:</b> Apply as a directed spray over the entire row surface at the rate of 2 pints per acre. Use up to 3 pints per acre in areas of heavy witchweed infestation. Use a minimum spray volume of 20 gallons per acre and a non-ionic surfactant at the rate of 2 pints per 100 gallons of spray.
<b>Postemergence</b>	1-2	<b>Repeat Applications:</b> In case of witchweed breakthrough a repeat application may be made at 1 to 2 pints per acre.

**Precautions:**

- Do not spray over the top of the corn, as this may result in severe corn injury. Spray should contact only the lower 3 to 8 inches of the corn stalk and any leaves in this zone. Spray droplets contacting the lower leaves will cause necrotic spotting or streaking of sprayed tissue.

**Crop-Specific Restrictions:**

- Do not apply more than 5 pints (1.25 lb active) of Willowood OxyFlo 2EC per acre during the growing season.
- Do not apply within 60 days of harvest.
- Do not use corn plants from a treated field for green chop, ensilage, forage, or fodder.

**COTTON**

**Application Methods and Equipment:** Willowood OxyFlo 2EC may be applied as a post-direct spray to cotton a minimum of 6 to 8 inches tall. Care must be exercised to avoid spray contact with the cotton leaves. Use rigid precision ground spray equipment and spray shields to prevent spray contact with cotton foliage. Use branch lifters or shields, as necessary, to avoid contact of directed sprays with cotton plant.

Accurate placement of spray nozzles is essential for uniform coverage of weeds and to minimize injury to cotton plants. Use a minimum broadcast spray volume of 20 gallons per acre and operate the sprayer at the minimum spray pressure recommended by the spray nozzle manufacturer. Willowood OxyFlo 2EC may be applied as a post-direct spray with only 2 flat fan nozzles per row (1 nozzle on each side of the row). For optimum coverage, use 4 flat fan nozzles per row (2 nozzles on each side of the row). The 2 forward nozzles should point forward and downward while the rear nozzles should point to the rear and downward. With either spraying setup, nozzles should be carefully adjusted to cover the weed foliage with minimum contact to cotton plants. Willowood OxyFlo 2EC may also be applied as a band application. **Do not use hollow cone nozzles.**

**Tank Mixing:** For control of additional broadleaf and grass weeds, Willowood OxyFlo 2EC may be applied as a postemergence directed spray in tank mix combination with other herbicides registered for postemergence use in cotton (see Tank Mixing Precautions under Mixing Directions).

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Postemergence</b>	1-2	<p>Apply as a post-directed spray. For optimum control, use the 2 pint per acre rate on actively growing weed seedlings with no more than 4 true leaves (not counting cotyledon leaves).</p> <p>Effective control of succulent weeds at the 2- to 3-leaf stage can usually be obtained at the 1 pint per acre rate. See Mixing Directions for surfactant recommendations.</p> <p>Where available, irrigation may be applied prior to application of Willowood OxyFlo 2EC to encourage maximum weed emergence.</p> <p>Irrigation following application will improve preemergence activity of Willowood OxyFlo 2EC against nightshade and groundcherry species.</p>

**Precautions:**

- Do not apply to cotton less than 6 inches tall or severe crop injury will result.
- Exercise care to avoid spray contact with cotton leaves. Leaves accidentally sprayed will exhibit necrotic (dead) spots and may be dropped from the plant. Crop injury may be enhanced if application is made when excessive soil moisture is present or rainfall occurs immediately after application, however, cotton will outgrow this condition and develop normally.

**Crop-Specific Restrictions:**

- **Western Cotton (AZ and CA):** Do not apply more than 2 pints (0.5 lb active) of Willowood OxyFlo 2EC per acre in a single application, or more than a total of 4 pints (1.0 lb active) of Willowood OxyFlo 2EC per broadcast acre per season as a result of multiple applications. Do not apply within 75 days of harvest.
- **Southern Cotton (All other states):** Do not apply more than 2 pints (0.5 lb active) of Willowood OxyFlo 2EC per acre per season as a result of a single application or multiple applications. Do not apply within 90 days of harvest.

**Key Weeds Controlled:****Postemergence**

cocklebur, common	nightshade, hairy
croton, tropic	pigweed, redroot
groundcherry, cutleaf	poinsettia, wild*
groundcherry, Wright	purslane, common
jimsonweed	sesbania, hemp
lambsquarters, common	sicklepod**
morningglory, annual (up to 6 leaf)	sida, prickly (teaweed)*
nightshade, American black	smartweed, Pennsylvania
nightshade, black	velvetleaf

\*Multiple applications may be required for acceptable control.

\*\*Post-direct applications of Willowood OxyFlo 2EC will control or suppress seedlings not exceeding the one true leaf stage.

### **COTTONWOOD**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	4-6	Willowood OxyFlo 2EC may be applied as a single or split application. Apply as a directed spray to soil at the base of cottonwood trees. Use the higher rate in the rate range for extended preemergence weed control or for postemergence control of weeds up to the 6 leaf stage. The addition of a non-ionic surfactant at 2 pints per 100 gallons of spray will enhance the postemergence activity of Willowood OxyFlo 2EC on emerged weeds.
<b>Precautions:</b> <ul style="list-style-type: none"><li>• Apply Willowood OxyFlo 2EC immediately after transplant only to dormant healthy cottonwood stock.</li><li>• In established stands, do not allow sprays of Willowood OxyFlo 2EC to contact cottonwood foliage. In newly established cottonwood plantings, use spray shields, if necessary, to prevent exposure of green bark and foliage.</li></ul>		
<b>Crop-Specific Restrictions:</b> <ul style="list-style-type: none"><li>• Do not apply more than 6 pints per acre of Willowood OxyFlo 2EC in a single application or more than 18 pints per acre per year.</li></ul>		

#### **Key Weeds Controlled:**

groundsel, common

knotweed, prostrate

lambsquarters, common

mustard, hedge

shepherdspurse

smartweed, Pennsylvania

## EUCALYPTUS

Apply Willowood OxyFlo 2EC for preemergence and postemergence control of listed broadleaf weeds in established eucalyptus plantings.

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b> <b>Postemergence</b>	4-6	<p><b>Directed Spray:</b> Willowood OxyFlo 2EC may be applied as a single or split application. Apply as a directed spray to soil at the base of eucalyptus trees.</p> <p>Use the higher rate in the range for extended preemergence weed control or for postemergence control of weeds up to the 6 leaf stage. The addition of a non-ionic surfactant at the rate of 2 pints per 100 gallons of spray will enhance the postemergence activity of Willowood OxyFlo 2EC on emerged weeds.</p> <p><b>Over-the-Top Application:</b> In new plantings, apply Willowood OxyFlo 2EC just before or immediately after transplanting eucalyptus seedlings that are in a dormant condition (i.e., leaves may be present, but terminal growth has hardened off and terminal buds have formed). In established plantings, Willowood OxyFlo 2EC may be applied as an over-the-top spray when plants are in a dormant condition.</p>
<p><b>Precautions:</b></p> <ul style="list-style-type: none"> <li>At transplant, apply Willowood OxyFlo 2EC only to healthy “dormant” eucalyptus stock. In established plantings, use spray shields, if needed, to prevent exposure of foliage and bark of small and/or actively growing plants.</li> <li>To avoid phytotoxicity, make over-the-top applications only to eucalyptus trees in a dormant condition. Do not make over-the-top applications after bud break and resumption of active growth.</li> </ul>		
<p><b>Crop-Specific Restrictions:</b></p> <ul style="list-style-type: none"> <li>Do not apply more than 6 pints of Willowood OxyFlo 2EC per acre in a single application or more than 18 pints per acre per year.</li> </ul>		

### Key Weeds Controlled:

#### Preemergence

burclover  
 cheeseweed (malva)  
 fiddleneck, coast  
 filaree, broadleaf  
 filaree, redstem  
 filaree, whitestem  
 groundsel, common  
 henbit  
 knotweed, prostrate  
 lambsquarters, common  
 lettuce, prickly  
 pigweed, redroot  
 purslane, common  
 redmaids  
 rocket, London  
 shepherdspurse  
 sowthistle, annual  
 spurge, prostrate  
 spurge, spotted

#### Postemergence

cheeseweed (malva)  
 fiddleneck, coast  
 filaree, broadleaf\*  
 filaree, redstem\*  
 filaree, whitestem\*  
 groundsel, common  
 henbit  
 minerslettuce  
 nettle, burning  
 pigweed, redroot  
 redmaids  
 shepherdspurse  
 sowthistle, annual

\* At the 6-pint rate, Willowood OxyFlo 2EC will provide control of filaree up to the 6-leaf stage.

## **USE ON FALLOW BEDS**

### **(Do not use prior to planting soybeans in California)**

Used alone or in a tank mix combination with glyphosate, Willowood OxyFlo 2EC provides preemergence and/or postemergence control of winter annual broadleaf weeds on land to be planted to crops.

Prior to planting, treated fallow beds should be thoroughly tilled (incorporated) to a depth of at least 2.5 inches. Willowood OxyFlo 2EC is no longer herbicidally effective once the active layer in the soil surface is disrupted by soil incorporation.

**Aerial Application:** Willowood OxyFlo 2EC may be aerially applied for weed control in fallow beds. Follow requirements for Aerial Application in the Product Information section of this label.

### **Minimum Treatment to Planting Intervals for listed crops:**

Direct Seeded Crops	Minimum Treatment-to-Planting Interval	
	Willowood OxyFlo 2EC (Up to 1 pint/acre)	Willowood OxyFlo 2EC (>1 to 2 pints/acre)
carrot	90 days	90 days
cotton	7 days	7 days
potato	60 days	60 days
sugar beet	60 days	90 days
other root/tuber crops	90 days	90 days
onions	180 days	180 days
other bulb vegetables	180 days	180 days
cabbage	90 days	90 days
cauliflower	90 days	90 days
other brassica crops	120 days	120 days
lettuce	90 days	120 days
other leafy vegetables (except brassica crops)	120 days	120 days
pepper	90 days	120 days
tomato	60 days	120 days
other fruiting vegetables	120 days	120 days
cantaloupe	60 days	90 days
squash	90 days	120 days
watermelon	60 days	60 days
other cucurbits	90 days	120 days
dry beans	60 days	60 days



Direct Seeded Crops	Minimum Treatment-to-Planting Interval	
	Willowood OxyFlo 2EC (Up to 1 pint/acre)	Willowood OxyFlo 2EC (>1 to 2 pints/acre)
peanut	60 days	60 days
other legume vegetables	60 days	60 days
safflower	60 days	60 days
soybeans (Except California)	7 days	7 days
cereal grains: Including barley, buckwheat, corn, proso, millet, pearl millet, oats, popcorn, rice, rye, sorghum, triticale, wheat, and wild rice	10 months	10 months
cotton and soybean	(See specified labeling for fallow beds to be planted to cotton or soybeans.)	

Transplanted Crops	Minimum Treatment-to-Planting Interval	
	Willowood OxyFlo 2EC (up to 1 pt/acre)	Willowood OxyFlo 2EC (>1 to 2 pints/acre)
celery	30 days	30 days
conifer	0 days	0 days
garlic	0 days	30 days
grape/kiwi	0 days	0 days
onion	0 days	30 days
pepper	30 days	30 days
strawberries	30 days	30 days
tomato	30 days	30 days
treefruit/nut/citrus	0 days	0 days

Weed Control	Rate (pt/acre)	Specific Use Directions
--------------	----------------	-------------------------

<b>Preemergence Postemergence</b>	1-2	<p>Use 20 or more gallons of spray volume per acre and increase spray volume for dense weed growth.</p> <p>Use the 1 pint per acre rate for up to 4 weeks of preemergence control and postemergence control of susceptible weeds up to the 4-leaf stage. Use the 2 pint per acre rate for up to 8 weeks of preemergence control and postemergence control of susceptible weeds up to the 6-leaf stage. Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks after application.</p> <p>A tank mix with glyphosate is recommended if the treatment area contains dense weed populations, oversized weed seedlings, volunteer grains, annual grasses or under unfavorable environmental conditions.</p> <p><b>Outside of California:</b> For enhanced contact activity (burndown/suppression) tank mix 6.5 fl oz of Willowood OxyFlo 2EC with the labeled rate of either glyphosate or paraquat (Gramoxone). Apply at the application rate and weed growth stages recommended in the respective tank mix product label.</p>
---------------------------------------	-----	--

**Precautions:**

- **Failure to achieve thorough and complete incorporation, or to follow the recommended treatment-planting interval, may result in stand reduction and/or vigor reduction of the planted crop.**
- Crop injury may be enhanced if newly seeded crops or transplants are under stress due to drought, flooding, excessive fertilizer or soil salts, low soil temperatures, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.
- **Exercise extreme care to avoid herbicide contact with any desirable dormant or non-dormant crop, plant, tree or vegetation as severe injury may result.**

**Crop-Specific Restrictions:**

- Do not apply more than 2 pints of Willowood OxyFlo 2EC per acre per fallow season.

**Key Weeds Controlled:** Willowood OxyFlo 2EC provides preemergence and postemergence control of the following weeds on fallow beds:\*

buttercup, smallflower	mustard species
cheeseweed (malva)	nettle, burning
eveningprimrose, cutleaf**	oxalis
fiddleneck, coast	pigweed, redroot
filaree, broadleaf	purslane, common
filaree, redstem	redmaids
geranium, Carolina	rocket, London
groundcherry, cutleaf	shepherdspurse
groundsel, common	sida, prickly
henbit	sowthistle, annual
ladysthumb	velvetleaf (wild cotton)
minerslettuce	

\*Thorough spray coverage is essential to maximize the postemergence activity of Willowood OxyFlo 2EC. For postemergence control when applied by air, a tank mixture of Willowood OxyFlo 2EC with either glyphosate or paraquat (Gramoxone) is recommended.

\*\*Requires maximum rate and/or multiple applications for effective control.

**FALLOW BED USE PRIOR TO TRANSPLANTING STRAWBERRIES OR PEPPERS GROWN IN PLASTIC CULTURE (CALIFORNIA ONLY)**

## Product Information

In California, apply this product broadcast or banded as a fallow bed application to pre-formed beds prior to planting strawberries or peppers grown in plastic culture. Use soil moisture to activate the product soon after application by irrigating the beds with 0.5 inch of sprinkler irrigation and then put plastic down anytime during the 30-day treatment-to-planting interval. If there is adequate soil moisture, apply plastic to the beds as soon as possible after application and allow the moisture which condenses and accumulates beneath the plastic to thoroughly wet the treated soil. Mechanical incorporation of the fallow-bed treatment prior to laying plastic is not required. Not disturbing the soil surface can allow for extended weed control. Not incorporating the soil surface increases the potential for crop injury, especially under wet conditions. Therefore, incorporate the treatment if the risk of crop injury is not acceptable. Follow directions for use and the minimum treatment-to-planting intervals outlined for fallow bed applications.

### **FALLOW BED USE PRIOR TO TRANSPLANTING PEPPERS, STRAWBERRIES OR TOMATOES GROWN IN PLASTIC CULTURE FLORIDA, GEORGIA, NORTH CAROLINA, SOUTH CAROLINA AND VIRGINIA ONLY**

## Product Information

In Florida, Georgia, North Carolina, South Carolina and Virginia, Willowood OxyFlor 2EC is effective as a preemergence broadcast or banded treatment to pre-formed beds as a fallow-bed application prior to planting of peppers, strawberries or tomatoes grown in plastic culture. Put down plastic anytime during the 30-day treatment-to-planting interval. Incorporation of the fallow-bed treatment prior to laying plastic is not required and can allow extended weed control. However, not incorporating increases the potential for crop injury, especially under wet conditions. Therefore, incorporate the treatment if the risk of crop injury is not acceptable. Follow directions for use and the minimum treatment-to-planting intervals outlined above for fallow bed applications.

### **FALLOW BEDS TO BE PLANTED TO COTTON AND SOYBEANS**

#### **Ground or Aerial Application of Willowood OxyFlo 2EC on Fallow Beds to Be Planted to Cotton or Soybeans**

#### **Not for Use on Fallow Beds to be Planted to Soybeans in California**

## Product Information

Willowood OxyFlo 2EC is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphogan) or paraquat (Parazone 3SL) for the control of winter annual broadleaf weeds in fallow beds to be planted to cotton or soybeans. Do not apply this product within 7 days prior to planting. Work the fallow beds thoroughly to a depth of at least 2 inches prior to planting. It is important to thoroughly break the soil surface prior to planting. Do not expect weed control following breaking of the soil surface.

EXERCISE EXTREME CARE TO AVOID HERBICIDE CONTACT WITH ANY DESIRABLE DORMANT OR NON-DORMANT CROP, PLANT, TREE, OR VEGETATION AS SEVERE INJURY MAY RESULT.

#### **Willowood OxyFlo 2EC Used Alone**

## Dosage

Apply this product at 1 to 2 pints (0.25 to 0.5 lbs. active) per broadcast acre. The lower rate (1 pint per acre) provides up to 4 weeks of preemergence control of susceptible weeds and provides postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre) provides preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible weeds (up to 6-leaf stage). Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

## Weeds Controlled

Willowood OxyFlo 2EC provides preemergence and postemergence\* control of the following weeds when used at specified dosages and weed stage:

Buttercup, Smallflower	Mustard Species
Cheeseweed (Malva)	Nettle, Burning
Eveningprimrose, Cutleaf**	Oxalis
Fiddleneck, Coast	Pigweed, Redroot
Filaree, Broadleaf	Purslane, Common
Filaree, Redstem	Redmaids
Geranium, Carolina	Rocket, London

Groundcherry, Cutleaf	Shepherdspurse
Groundsel, Common	Sida, Prickly
Henbit	Sowthistle, Annual
Ladysthumb	Velvetleaf (Wild Cotton)
Miner's Lettuce	

\*Thorough spray coverage is essential to maximize the postemergence activity of this product. For postemergence control when applied by air, use a tank mixture of Willowood OxyFlo 2EC with either glyphosate (Glyphogan) or paraquat (Parazone 3SL).

\*\*Requires maximum rate and/or multiple applications for effective control.

### **Tank Mixes with Willowood OxyFlo 2EC**

**IMPORTANT:** Read and observe all label directions before using. When tank mixing, always read all individual manufacturer's labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

#### **Dosage:**

Willowood OxyFlo 2EC can be tank-mixed with either glyphosate (Glyphogan) or paraquat (Parazone 3SL) to obtain postemergence control of annual grassy weeds, volunteer grains, and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lbs. active) of this product with labeled rates of either glyphosate (Glyphogan) or paraquat (Parazone 3SL). Apply at the specified rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

**Outside of California:** For enhanced contact activity (burndown/suppression) to either glyphosate (Glyphogan) or paraquat (Parazone 3SL), add this product at a rate of 6.5 ounces (0.1 lb. active) per acre to labeled rates of either glyphosate (Glyphogan) or paraquat (Parazone 3SL). Apply at the specified rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

If a fallow bed treatment is applied thirty days or more prior to planting and at least three significant rainfalls (0.25 inch or greater) have occurred following application, cotton or soybeans can be planted directly into the stale seedbed. If these conditions cannot be met, soil incorporation is required as directed above.

### **Method of Application**

#### **Ground Application**

Apply Willowood OxyFlo 2EC in a minimum of 20 gallons of water per acre. Increase the volume of water used as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Calibrate spray equipment carefully before each use.

#### **Aerial Application**

Apply this product using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 10 gallons per acre (in California, minimum 10 GPA when applied alone or tank mixed with paraquat (Parazone 3SL)). Make applications at a height of 6 to 10 feet above the soil surface. Do not place the nozzles on the spray booms any closer to the wing or rotor tips than 3/4 of the span; this will minimize the formation of spray or wing tip vortices roll. Nozzles must be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100 microns or less in diameter.

#### **Important**

Aerial applicators must be familiar with this label and follow the use precautions. Spraying Willowood OxyFlo 2EC in a manner other than as specified is done at the user's risk. Users are responsible for all loss or damage that result from such spraying. In addition, aerial applicators must follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, apply the most restrictive situations to avoid drift hazards.

The use of a drift control agent may be required by local regulations. However, the drift control agent may decrease the weed control effectiveness.

### **Fallow Bed (Cotton, Soybeans)**

#### **Specified Use Restrictions**

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not apply more than 2 pints (0.5 lbs. active) of this product per acre per fallow season.
- Do not apply this product within 7 days prior to planting of cotton or soybeans.

## **FALLOW BEDS TO BE PLANTED TO FIELD CORN**

**(ARKANSAS, LOUISIANA AND MISSISSIPPI ONLY)**

### **Ground or Aerial Application of Willowood OxyFlo 2EC on Fallow Beds to be Planted to Field Corn**

**Use only on Fallow Beds to be Planted to Field Corn in Arkansas, Louisiana and Mississippi**

#### **Product Information**

Willowood OxyFlo 2EC is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphogan) or paraquat (Parazone 3SL) for the control of winter annual broadleaf weeds in fallow beds to be planted to corn. Do not apply this product within 7 days prior to planting. Unless otherwise specified in this label, work the fallow beds thoroughly to a depth of at least 2 inches prior to planting. It is important to thoroughly break the soil prior to planting. Do not expect weed control following breaking of the soil surface.

If a fallow bed treatment is applied thirty days or more prior to planting and at least three significant rainfalls (0.25 inch or greater) have occurred following application, corn can be planted directly into the stale seedbed. If these conditions cannot be met, soil incorporation is required as directed above.

EXERCISE EXTREME CARE TO AVOID HERBICIDE CONTACT WITH ANY DESIRABLE DORMANT OR NON-DORMANT CROP, PLANT, TREE, OR VEGETATION AS SEVERE INJURY MAY RESULT.

#### **Willowood OxyFlo 2EC Used Alone**

##### **Dosage**

Apply this product at 1 to 2 pints (0.25 to 0.5 lbs. active) per broadcast acre. The lower rate (1 pint per acre) provides up to 4 weeks of preemergence control of susceptible weeds and provides postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre) provides preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible weeds (up to 6-leaf stage). Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

##### **Weeds Controlled**

This product provides preemergence and postemergence\* control of the following weeds when used at specified dosages and weed stage:

Buttercup, Smallflower	Mustard Species
Cheeseweed (Malva)	Nettle, Burning
Eveningprimrose, Cutleaf**	Oxalis
Fiddleneck, Coast	Pigweed, Redroot
Filaree, Broadleaf	Purslane, Common
Filaree, Redstem	Redmaids
Geranium, Carolina	Rocket, London
Groundcherry, Cutleaf	Shepherdspurse
Groundsel, Common	Sida, Prickly
Henbit	Sowthistle, Annual
Ladysthumb	Velvetleaf (Wild Cotton)
Miner's Lettuce	

\*Thorough spray coverage is essential to maximize the postemergence activity of Willowood OxyFlo 2EC. For postemergence control when applied by air, use a tank mixture of this product with either glyphosate (Glyphogan) or paraquat (Parazone 3SL).

\*\*Requires maximum rate and/or multiple applications for effective control.

## **Tank Mixes With Willowood OxyFlo 2EC**

**IMPORTANT:** Read and observe all label directions before using. When tank mixing, always read all individual manufacturer's labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

### **Dosage:**

This product can be tank-mixed with either glyphosate (Glyphogan) or paraquat (Parazone 3SL) to obtain postemergence control of annual grassy weeds, volunteer grains, and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lbs. active) of this product with labeled rates of either glyphosate (Glyphogan) or paraquat (Parazone 3SL). Apply at the specified rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

For enhanced contact activity (burndown/suppression) to either glyphosate (Glyphogan) or paraquat (Parazone 3SL), add Willowood OxyFlo 2EC at a rate of 3.5 to 7 fl. oz. (0.05 to 0.1 lb. active) per acre to labeled rates of either glyphosate (Glyphogan) or paraquat (Parazone 3SL). Apply at the specified rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

### **Method of Application**

#### **Ground Application**

Apply in a minimum of 20 gallons of water per acre. Increase the volume of water used as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Calibrate spray equipment carefully before each use.

#### **Aerial Application**

Apply this product using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 5 gallons per acre. Make applications at a height of 6 to 10 feet above the soil surface. Do not place the nozzles on the spray booms any closer to the wing or rotor tips than 3/4 of the span; this will minimize the formation of spray or wing tip vortice roll. Nozzles must be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100 microns or less in diameter.

### **Important**

Aerial applicators must be familiar with this label and follow the use precautions. Spraying Willowood OxyFlo 2EC in a manner other than as specified is done at the user's risk. Users are responsible for all loss or damage that result from such spraying. In addition, aerial applicators must follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, apply the most restrictive situations to avoid drift hazards.

## **Fallow Bed (Field Corn) – Arkansas, Louisiana and Mississippi**

### **Specific Use Restrictions**

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not apply more than 2 pints (0.5 lb. active) of this product per acre per fallow season.
- Do not apply this product within 7 days prior to planting of corn.
- Do not use corn plants from a treated field for green chop, ensilage, forage or fodder.
- Do not feed or allow animals to graze on any areas treated with this product.
- Do not treat ditch banks or waterways with this product.
- Chemigation: Do not apply this product through any type of irrigation system except as specified on other approved supplemental labeling.

**IMPORTANT:** Treated soil must be thoroughly incorporated to a depth of 4 inches after harvest (or abandoning) of the treated crop but prior to planting of the rotational crop. Failure to achieve this thorough and complete incorporation or to follow the required minimum plant back interval may result in crop injury, stand reduction, and/or vigor reduction of the plant-back crop. See specific fallow bed labeling regarding crop planting information for applications of this product made to a fallow bed or fallow field.

## **FALLOW BEDS TO BE PLANTED TO FIELD CORN**

### **(CALIFORNIA ONLY)**

#### **Ground or Aerial Application of Willowood OxyFlo 2EC on Fallow Beds to be Planted to Field Corn**

#### **Use Only on Fallow Beds to be Planted to Field Corn in California**

## Product Information

This product is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with glyphosate (Glyphogan) or paraquat (Parazone 3SL) for the control of winter annual broadleaf weeds in fallow beds to be planted to corn. Do not apply this product within 60 days prior to planting. Work the fallow beds thoroughly to a depth of at least 2.5 inches prior to planting. It is important to thoroughly break the soil surface prior to planting. Do not expect weed control following breaking of the soil surface.

If a fallow bed treatment is applied sixty days or more prior to planting and at least three significant rainfalls (0.25 inch or greater) have occurred following application, corn can be planted directly into the stale seedbed. If these conditions cannot be met, soil incorporation is required as directed above.

EXERCISE EXTREME CARE TO AVOID HERBICIDE CONTACT WITH ANY DESIRABLE DORMANT OR NON-DORMANT CROP, PLANT, TREE, OR VEGETATION AS SEVERE INJURY MAY RESULT.

### Willowood OxyFlo 2EC Used Alone

#### Dosage

Apply this product at 1 to 2 pints (0.25 to 0.5 lb. active) per broadcast acre. The lower rate (1 pint per acre) provides up to 4 weeks of preemergence control of susceptible weeds and provides postemergence control of susceptible weeds (up to 4-leaf stage). The higher rate (2 pints per acre) provides preemergence control of susceptible weeds for up to 8 weeks and postemergence control of susceptible weeds (up to 6-leaf stage). Best preemergence control is achieved when irrigation or rainfall occurs within 3 or 4 weeks following application.

#### Weeds Controlled

This product provides preemergence and postemergence\* control of the following weeds when used at specified dosages and weed stage:

Buttercup, Smallflower	Mustard Species
Cheeseweed (Malva)	Nettle, Burning
Eveningprimrose, Cutleaf**	Oxalis
Fiddleneck, Coast	Pigweed, Redroot
Filaree, Broadleaf	Purslane, Common
Filaree, Redstem	Redmaids
Geranium, Carolina	Rocket, London
Groundcherry, Cutleaf	Shepherdspurse
Groundsel, Common	Sida, Prickly
Henbit	Sowthistle, Annual
Ladysthumb	Velvetleaf (Wild Cotton)
Miner's Lettuce	

\*Thorough spray coverage is essential to maximize the postemergence activity of Willowood OxyFlo 2EC. For postemergence control when applied by air, use a tank mixture of this product with either glyphosate (Glyphogan) or paraquat (Parazone 3SL).

\*\*Requires maximum rate and/or multiple applications for effective control.

### Tank Mixes With Willowood OxyFlo 2EC

**IMPORTANT:** Read and observe all label directions before using. When tank mixing, always read all individual manufacturer's labels. In interpreting all labels for the tank mix, the most restrictive situations must apply.

#### Dosage:

This product can be tank-mixed with either glyphosate (Glyphogan) or paraquat (Parazone 3SL) to obtain postemergence control of annual grassy weeds, volunteer grains, and broadleaf weeds. Tank mix 1 to 2 pints (0.25 to 0.5 lb. active) of this product with labeled rates of either glyphosate (Glyphogan) or paraquat (Parazone 3SL). Apply at the specified rates and growth stages to susceptible weed species in a manner consistent with the respective labels.

#### Method of Application

##### Ground Application

Apply in a minimum of 20 gallons of water per acre. Increase the volume of water used as the weeds become taller and more dense. Use a low-pressure sprayer equipped with flat fan nozzles. Calibrate spray equipment carefully before each

use.

### **Aerial Application**

Apply this product using swirl jet or hollow cone nozzles and a spray pressure less than 40 psi to deliver a minimum spray volume of 5 gallons per acre. Make applications at a height of 6 to 10 feet above the soil surface. Do not place the nozzles on the spray booms any closer to the wing or rotor tips than 3/4 of the span; this will minimize the formation of spray or wing tip vortex roll. Nozzles must be spaced and positioned to produce a uniform spray pattern and to minimize or eliminate the formation of droplets 100 microns or less in diameter.

### **Important**

Aerial applicators must be familiar with this label and follow the use precautions. Spraying this product in a manner other than as specified is done at the user's risk. Users are responsible for all loss or damage that result from such spraying. In addition, aerial applicators must follow all applicable state and local regulations and ordinances. In interpreting the label and local regulations, apply the most restrictive situations to avoid drift hazards.

Crop injury may be enhanced if newly seeded crops are under stress due to drought, flooding, excessive fertilizer or soil salts, low soil temperatures, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.

### **Fallow Bed (Corn) – California**

#### **Specific Use Restrictions**

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not apply more than 2 pints (0.5 lb. active) of Willowood OxyFlo 2EC per acre per fallow season. Do not apply more than 0.5 lb. of oxyfluorfen active ingredient per acre per fallow season as a result of single or multiple applications of this or other oxyfluorfen formulations.
- Do not apply this product within 60 days prior to planting of corn.
- Before planting field corn, treated soil must be thoroughly mixed to a depth of at least 2.5 inches.
- Chemigation: Do not apply this product through any type of irrigation system except as specified elsewhere on this label.
- Do not use on sweet corn.
- Do not use corn plants from a field treated with Willowood OxyFlo 2EC for green chop, ensilage, forage or fodder.
- Do not feed or allow animals to graze on any areas treated with this product.
- Application may be made in a minimum of 20 gals. of water/A using ground equipment or 5 gals. of water/A by air. Applications may be made alone or as a tank mix with other herbicides.
- Do not apply more than 0.5 lb. active ingredient per year.



## **FALLOW LAND**

### **(For Use Only in Idaho, Oregon and Washington)**

Used alone or in a tank mix combination with glyphosate, Willowood OxyFlo 2EC provides preemergence and/or postemergence control of listed annual broadleaf weeds in a fallow land system. Willowood OxyFlo 2EC may be used to reduce weed growth prior to the establishment of dry soil mulch. Use is restricted to summer fallow on land that will be planted the following year to winter wheat, barley or oats.

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	0.5-2	<b>Willowood OxyFlo 2EC Alone:</b> Preemergence weed control occurs as seedling weeds come in contact with the soil-applied herbicide during emergence. Postemergence weed control is most effective when Willowood OxyFlo 2EC is applied to seedling weeds less than 4 inches in height. Apply Willowood OxyFlo 2EC in 15 or more gallons of water per acre and increase spray volume if weed growth is dense. Use of an 80% active nonionic surfactant cleared for use on growing crops is recommended for optimum postemergence weed control.
<b>Tank Mixing:</b> For postemergence control of annual grass weeds, 0.5 – 2 pts/acre of Willowood OxyFlo 2EC may be tank mixed with labeled rates of glyphosate. Follow label instructions for Fallow and Reduced Tillage Systems for the glyphosate product. Refer to Mixing Directions section for Tank Mixing Precautions.		
<b>Use Restrictions for Fallow Land:</b> <ul style="list-style-type: none"><li>• Do not apply more than 2 pints per acre per application or more than 2 pints per use season.</li></ul>		

**Key Weeds Controlled:** Willowood OxyFlo 2EC provides preemergence and postemergence control of the following weeds on fallow land.

fiddleneck, coast

henbit

lettuce, prickly (china lettuce)

mustard, blue (purple mustard)

mustard, tumble (Jim hill mustard)

pigweed, redroot

purslane, common

shepherdspurse

sowthistle, annual

## **GARBANZO BEANS**

**(For Use Only in Arizona and California)**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence</b>	1	Apply after planting but prior to weed or crop emergence as a single broadcast application using a spray volume of 20 or more gallons of water per acre.
<b>Precautions:</b> <ul style="list-style-type: none"><li>• Garbanzo beans are tolerant to preemergence application of Willowood OxyFlo 2EC, however, under certain conditions, severe but temporary crop injury may occur. A heavy splashing rain shortly after crop emergence or wet soil conditions during early growth stages can cause leaf cupping, crinkling, stunting or defoliation of the garbanzo seedlings. Injury, when it occurs, is usually limited to the first few leaves that develop after plants emerge from the soil. Delays in crop development and/or maturity may result, but Garbanzo beans do recover with little to no impact on yield.</li></ul>		
<b>Crop-Specific Restrictions:</b> <ul style="list-style-type: none"><li>• Do not apply more than 1 pint per acre of Willowood OxyFlo 2EC in a single application.</li><li>• Do not use bean vines for livestock feed or hay.</li></ul>		

### **Key Weeds Controlled:**

#### **Preemergence**

groundsel, common  
mallow, little  
rocket, London  
shepherdspurse

## **GARLIC**

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

**Cultural Considerations:** For optimum preemergence weed control, the soil surface should be smooth and free of excessive trash (clippings, plant residues, etc.). Following application, treated beds should be left undisturbed during the time period for which weed control is desired. Cultural practices that result in soil disturbance or redistribution or untreated soil can result in reduced weed control.

#### **Direct Seeded Garlic (Postemergence Application):**

<b>Weed Control</b>	<b>Rate (per acre)</b>	<b>Specific Use Directions</b>
---------------------	------------------------	--------------------------------

<b>Postemergence</b>	2-4 fl oz	<b>Northeastern States Including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont:</b> Apply Willowood OxyFlo 2EC at 2 to 4 fl oz per acre to seeded garlic that has at least 3 true leaves using ground equipment. Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.
<b>Postemergence</b>	0.5-1 pt	<b>Western States Including Arizona, Colorado, Idaho, Nevada, New Mexico, Oregon, Texas, Utah and Washington:</b> Apply Willowood OxyFlo 2EC at 0.5 to 1 pt per acre to seeded garlic that has at least 2 true leaves using ground equipment. Multiple treatments at 0.5 to 1 pt per acre may be applied up to a maximum of 2.0 pints per acre per use season. For optimum postemergence weed control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf stage may result in reduced weed control.
<b>Postemergence</b>	0.5 pt	<b>All Other States:</b> Apply Willowood OxyFlo 2EC at 0.5 pt per acre to seeded garlic that has at least 2 true leaves using ground equipment. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.

<b>Direct Seeded Garlic (California Only)</b>		
<b>Weed Control</b>	<b>Rate (per/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	1 pt	<p><b>Application after planting but Prior to Garlic Emergence:</b> Apply Willowood OxyFlo 2EC after planting, but prior to crop emergence, for preemergence control of listed broadleaf and grass weeds using ground, air or sprinkler irrigation (chemigation).</p> <p><b>Aerial Application:</b> Apply in a minimum spray volume of 10 gallons per acre. Follow Aerial Application instructions and precautions in the Product Information section of this label.</p> <p><b>Postemergence and Directed Application:</b> Apply Willowood OxyFlo 2EC as a directed over-the-top spray to garlic that is at least 12 inches tall. Accurate, uniform placement of directed postemergence sprays is essential for effective weed control and to minimize injury to garlic. Use low-pressure sprays and a minimum spray volume of 20 gallons per acre. Adjust nozzles for minimum spray contact with garlic plants, directing the spray to the soil at the base of garlic plants and adjacent bed top and furrow area. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing. Application at later than 4-leaf growth stage may result in reduced weed control.</p> <p><b>Sprinkler Irrigation (Portable Lateral or Solid Set)</b> <b>Preemergence or Postemergence:</b> Apply Willowood OxyFlo 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Apply after planting but prior to garlic emergence or postemergence when garlic is at least 12 inches tall. Follow the application directions and precautions for "Sprinkler Chemigation" given in the Chemigation section of this label.</p>

**Precautions:**

- **Garlic Response to Preemergence Applications of Willowood OxyFlo 2EC:** Following a preemergence application of Willowood OxyFlo 2EC, a chlorotic band around some of the leaves may be observed after the first irrigation (or rainfall) following garlic emergence.

<b>Transplanted Garlic: Postemergence Application Immediately after Planting</b>		
<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Postemergence</b>	up to 2 pt	<b>All States Except Northeastern States:</b> Transplanted garlic is most tolerant of a postemergence application immediately after transplanting. An application of up to 2 pints per acre may be made within two days after transplanting. If less than 2 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints per acre of Willowood OxyFlo 2EC per season as a result of multiple applications.
<b>Postemergence</b>	2-4 fl oz	<b>Northeastern States, including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont:</b> Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season.

**Key Weeds Controlled:**

canarygrass (annual)  
 evening primrose, cutleaf  
 groundsel, common  
 mallow, little (malva)  
 nightshade, black  
 pigweed, prostrate\*  
 pigweed, redroot\*

puncturevine  
 purslane, common\*  
 rocket, London  
 sage, lanceleaf  
 shepherdspurse\*  
 sowthistle, annual

\*Key weeds controlled at recommended rates in Northeastern States.

**Garlic – Crop-Specific Precaution (Postemergence Application)**

- Postemergence applications of Willowood OxyFlo 2EC may cause chlorotic leaf banding, necrotic lesions, or stunting of the garlic plants. Symptoms may be more severe if garlic emerged under cool, wet, overcast, or foggy weather. These conditions are temporary and should not affect the vigor or development of garlic plants.

**Garlic – Crop-Specific Restrictions (Applicable to All Methods of Application):**

- In all states except Northeastern states, do not apply until direct seeded garlic plants have two (2) fully developed true leaves. In the Northeastern states, do not apply until direct seeded garlic plants have three (3) fully developed true leaves. Application made prior to the specified growth stage may result in serious crop injury and is not recommended.
- Do not apply more than a total of 2 pints per acre of Willowood OxyFlo 2EC per use season as a result of multiple applications.
- Do not apply within 60 days of harvest.
- In direct seeded garlic (except in California), do not apply Willowood OxyFlo 2EC as a preemergence treatment.
- Use only on dry bulb garlic.
- Do not apply to garlic grown for seed.
- For weed control in garlic, do not mix Willowood OxyFlo 2EC with oils, surfactants, liquid fertilizers or pesticides except as specified on approved Willowood, LLC labeling.
- Do not apply to garlic plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases.

## **GRASSES GROWN FOR SEED**

**(For Use Only in Oregon and Washington)**

### **Product Information**

Use this product for late preemergence through early postemergence control/suppression of annual broadleaf weeds, annual grasses, and the seedling stage of perennial grasses, including volunteer seedlings, in established perennial grasses grown for seed. A minimum of 12 months of good growing conditions is required from the time of planting new stands of perennial grasses or until the perennial grasses are sufficiently well established (a minimum of 6 tillers or more) to be treated with this product.

**Note:** Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, do not apply this product directly to native prairie habitats. Maintain dosage rates at the lower end of the use rate range if effective seedling control can be achieved in an effort to lessen potential impacts to endangered species.

### **Crop Tolerance**

The application of this product to established perennial grasses will result in a chlorosis (yellowing) of the vegetation within two weeks after treatment. Some crop response symptoms may be present for up to three months following application. The use of this product can also result in a substantial reduction in vegetative growth by perennial grasses during the winter. Leaf chlorosis and reduction of vegetative growth is a typical and normal response, and seed yield of healthy, vigorous perennial grasses is typically not affected by fall application of this product. The grower accepts that conditions under which seed yield may be reduced are not fully understood and that a reduction in seed yield may occur. **DO NOT GRAZE FIELDS THAT HAVE BEEN TREATED WITH WILLOWOOD OXYFLO 2EC AS ILLEGAL RESIDUES MAY BE PRESENT IN THE VEGETATIVE FORAGE.** Grazing may also magnify crop injury and reduce the seed yield.

Crop tolerance to this product can be improved by limiting the amount of leaf tissue present on the established perennial grasses at time of application by such methods as propane flaming, intensive mechanical clipping (crew cutting), or livestock grazing prior to application.

### **Timing**

Early treatment is important for control of seedling grasses. Applications of this product at the onset of grass seed germination during the initial fall rains or fall sprinkler irrigation (late preemergence) or at the one-leaf growth stage (early postemergence) can provide somewhat better control of volunteer crop seedlings than applications at the two-leaf stage. Ample soil moisture soon after application is required for optimum performance against seedling grasses.

Applications to seedling grasses between the two- and six-leaf stage can result in partial control, varying with species. Single applications made to seedlings between the two- and six-leaf growth stages will cause injury and stunting, but regrowth will occur. If seedlings have not died within three to four weeks after treatment with this product and healthy green regrowth is visible, a second application may be needed. Willowood OxyFlo 2EC will not control established perennial grasses or seedlings of most annual and perennial grasses beyond the six-leaf stage.

### **Method of Application**

Apply as a broadcast application in a minimum spray volume of 20 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles at a minimum spray pressure of 30 psi. Do not exceed maximum spray

pressure of 60 psi. Calibrate spray equipment carefully before each use. Add a nonionic surfactant containing 80% active ingredient cleared for growing crops with this product at the rate of 0.12 to 0.5% spray volume for improved control of emerged seedlings. Maintain agitation until spraying is complete.

Tank mixtures of this product with other registered herbicides can result in enhanced crop injury. If a tank mixture is to be applied, make applications only to healthy, vigorous stands of perennial grasses. The decision to apply a tank mixture containing Willowood OxyFlo 2EC is at the sole discretion of the grower at the grower's risk.

## Dosage

Use this product on the following crops at the rates given:

**Kentucky bluegrass, tall fescue, orchardgrass, bentgrass, and perennial ryegrass:** In Oregon, apply an initial application of this product at a rate of 0.5 to 1.5 pints (0.125 to 0.375 lb. active) per acre. Do not apply more than 1.5 pints (0.375 lb. active) of this product per acre as a result of two applications in a single growing season. Make the initial application before the weed (or grass) seedlings to be controlled exceed the two-leaf stage (typically prior to December 15). A second application may be required for optimal control/suppression. For increased control of volunteer ryegrass seedlings in perennial ryegrass, do not apply more than 1.2 lbs. active of diuron (Diuron 4L) per acre in tank mixture with this product per season.

**In Washington,** apply a maximum of 1.5 pints (0.375 lb. active) per acre of this product after harvest. A single application of 0.5 to 1.5 pints (0.125 to 0.375 lb. active) per acre can be made. The maximum rate of 1.5 pints (0.375 lb. active) per acre of this product can be split and the initial application applied before the weed (or grass) seedlings exceed the two-leaf growth stage and no later than December 15. The final application must be completed prior to January 15.

**Fine fescues (Chewings, creeping red, and hard types):** In Oregon, apply a single application of this product at 0.5 pint (0.125 lb. active) per acre per season. Apply before the weed seedlings to be controlled exceed the two-leaf growth stage (Use Period: September 1 to December 15).

## Weeds Controlled

This product will provide control or suppression of the following weeds and volunteer crops when applied between the onset of germination and two-leaf growth stage:

Bentgrass	Fescue, Hard
Bluegrass, Annual	Fescue, Rattail
Bluegrass, Kentucky	Fescue, Tall
Bluegrass, Roughstalk	Orchardgrass
Brome, California (mountain)*	Ryegrass, Italian
Fescue, Fine (creeping red and Chewings)	Ryegrass, Perennial

\*These species are suppressed but not fully controlled by this product.

This product will also control or suppress many annual broadleaf weeds. Willowood OxyFlo 2EC will not control/suppress established perennial grasses or seedlings of most annual and perennial grasses beyond the 6-leaf stage.

## Grasses Grown for Seed (Oregon & Washington Only)

### Specific Use Restrictions

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Do not graze livestock in treated fields within 150 days of application.
- Do not apply this product within 150 days of harvest.
- Apply this product only by ground application equipment.
- Chemigation: Do not apply this product through any type of irrigation system.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. This product is phytotoxic to plant foliage.

- Do not treat ditch banks or waterways with this product.
- A 25 ft. vegetative buffer strip must be maintained between all areas treated with this product and lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.
- In Washington only: This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife, and fish. This product must not be used under this label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult other sections of this label for additional restrictions and precautions to protect aquatic organisms.

## **GRASSES GROWN FOR SEED**

### **PERENNIAL RYEGRASS AND TALL FESCUE GROWN FOR SEED**

**(For Use in Oregon Only)**

#### **Product Information**

Use this product for early postemergence suppression/control of various annual broadleaf weed seedlings in new plantings of fall seeded perennial ryegrass and tall fescue that have at least one to two tillers. Application to seedling plants that have not yet been tillered can result in severe crop injury or stand loss (plant death).

**Note:** Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, do not apply this product directly to native prairie habitats. Maintain dosage rates at the lower end of the use rate range if effective seedling control can be achieved in an effort to lessen potential impacts to endangered species.

#### **Crop Tolerance**

The application of this product to fall seeded perennial ryegrass and tall fescue (that have at least one to two tillers) will result in a chlorosis (yellowing) of the vegetation within two weeks after treatment. Some crop response symptoms may be present for up to three months following application. The use of this product can also result in a substantial reduction in vegetative growth by perennial grasses during the winter. Leaf chlorosis and reduction of vegetative growth is a typical and normal response, and seed yield of healthy, vigorous perennial grasses is typically not affected by fall application of this product. The grower accepts that conditions under which seed yield may be reduced are not fully understood and that a reduction in seed yield may occur. **DO NOT GRAZE FIELDS WITHIN 150 DAYS OF TREATMENT WITH THIS PRODUCT AS ILLEGAL RESIDUES MAY BE PRESENT IN THE VEGETATIVE FORAGE.** Grazing may also magnify crop injury and reduce the seed yield.

#### **Method of Application**

Apply as a broadcast application in a minimum spray volume of 20 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles at a minimum spray pressure of 30 psi. Do not exceed maximum spray pressure of 60 psi. Calibrate spray equipment carefully before each use. Add a nonionic surfactant containing 80% active ingredient cleared for growing crops with this product at the rate of 0.12 to 0.5% spray volume for improved control of emerged seedlings. Maintain agitation until spraying is complete.

Tank mixtures of this product with Nortron 4SC can result in enhanced crop injury. If a tank mixture is to be applied, make applications only to healthy, vigorous stands of perennial grasses. The decision to apply a tank mixture containing Willowood OxyFlo 2EC is at the sole discretion of the grower at the grower's risk.

### **Dosage**

Apply a single application of this product at a rate of 2 to 3 fl. oz. (0.025 to 0.375 lb. active) per acre, either alone or tank mixed with up to 3 pints of Nortron 4SC per acre. Some temporary crop response may occur, but typically it is only a transient effect and should not adversely impact yield. **DO NOT APPLY TO NEWLY PLANTED STANDS THAT ARE UNDER STRESS FROM ANY CAUSE AS THERE IS AN ENHANCED OPPORTUNITY FOR CROP INJURY TO OCCUR.** Control from the Willowood OxyFlo 2EC application is primarily directed at emerged seedling broadleaf weeds such as speedwell and groundsel, but control or suppression of other species is possible if tank mixed with Nortron 4SC. Overlaps (2X applications) will cause significant crop injury but should not result in excessive stand losses if the crop plants are at least one to two tillers when the applications are made.

### **Weeds Controlled**

This product will provide control or suppression of the following weeds and volunteer crops when applied between the onset of germination and two-leaf seedling growth stage:

Groundsel, common                      Speedwell

This product will also control or suppress many other annual broadleaf weeds.

### **Perennial Ryegrass and Tall Fescue – Oregon Only**

#### **Fall Seeded New Plantings Only**

#### **Specific Use Restrictions**

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Do not graze livestock in treated fields within 150 days of application.
- Do not apply this product within 150 days of harvest.
- Apply this product only by ground application equipment.
- Chemigation: Do not apply this product through any type of irrigation system.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. This product is phytotoxic to plant foliage.
- Do not treat ditch banks or waterways with this product.

#### **GUAVA (Bearing and Non-bearing)**

**(For Use Only in Hawaii)**



Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b>	5 – 8	<p><b>Preemergence or Postemergence:</b> In established guava plantings, apply preemergence or postemergence to weeds. Increase the spray volume to ensure adequate coverage in high densities of emerged weeds or heavy trash. Minimize contact with guava plants by directing the spray to the soil surface. Spray shields are suggested to minimize spray contact in young plantings.</p> <p>For broader spectrum postemergence control of grass and broadleaf weeds, Willowood OxyFlo 2EC may be applied in a tank mix combination with paraquat (Gramoxone) or glyphosate. Follow applicable use directions, precautions and limitations on the labels of the respective tank mix products.</p>
<b>Postemergence</b>	2 – 8	

**Precautions:**

- Prevent direct spray or drift from contacting green stems, fruit or foliage, as injury may result.
- Alone or in tank mix combination, Willowood OxyFlo 2EC should be applied to only healthy growing trees.
- Application of Willowood OxyFlo 2EC should be made only after new foliage growth has hardened off.

**Crop-Specific Restrictions:**

- Do not apply more than 8 pints per acre of Willowood OxyFlo 2EC in a single application or more than 16 pints per season.
- Do not apply Willowood OxyFlo 2EC within 1 day of harvest.

**Key Weeds Controlled:**

**Preemergence**

ageratum  
 buttonweed  
 crotalaria  
 purslane, common  
 spurge, garden

**Postemergence**

purslane, common  
 spurge, garden

**HORSERADISH**

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Weed Control	Rate (pt/acre)	Specific Use Directions
--------------	----------------	-------------------------

<b>Preemergence</b>	2	Apply Willowood OxyFlo 2EC after the horseradish roots have been planted but prior to emergence of new horseradish leaves. Emerged leaves that receive direct or indirect spray (drift) contact will be injured. If necessary, cultivate before application to destroy germinated weeds.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Do not apply Willowood OxyFlo 2EC to horseradish plantings that have been weakened or stressed due to unfavorable temperature conditions, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.</li> </ul>		
<b>Crop-Specific Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 2 pints of Willowood OxyFlo 2EC per acre per crop.</li> </ul>		

**Key Weeds Controlled:**

lambquarters, common pigweed, redroot purslane, common	shepherdspurse smartweed, pennsylvania
--	---

**JOJOBA**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	4 – 6	Initial application may be made when jojoba plants have reached a height of 6 inches or more. Use sufficient spray volume to ensure thorough coverage of dense weed growth. Sprays should be directed to the base of jojoba plants to avoid possible phytotoxicity to foliage. Spray shields are suggested for use in young plantings. Use higher rate in rate range for extended residual preemergence weed control. Make follow-up applications as necessary to maintain weed control. For early postemergence control of susceptible seedling weeds (less than 8 inches tall) apply Willowood OxyFlo 2EC at the rate of 4 pints per acre. Willowood OxyFlo 2EC may be applied at the rate of 6 pints per acre for postemergence control of weeds up to 12 inches tall. For optimum residual control, apply during the fall or winter months. Control may be unsatisfactory for weeds greater than 12 inches tall.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Avoid direct spray or drift contact with jojoba flowers or buds as severe injury may result.</li> <li>Over-the-top applications may cause burning, crinkling or bronzing of jojoba foliage, particularly to the youngest leaves, flowers, or buds present at the time of application.</li> </ul>		
<b>Crop-Specific Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 6 pints per acre per year.</li> </ul>		

**Key Weeds Controlled:**

**Preemergence**

**Postemergence**

burclover  
 fiddleneck, coast  
 filaree, broadleaf  
 filaree, redstem  
 filaree, whitestem  
 groundsel, common  
 henbit  
 knotweed, prostrate  
 lambsquarters, common  
 lettuce, prickly  
 mallow, little (malva, cheeseweed)  
 pigweed, redroot  
 purslane, common  
 redmaids  
 rocket, London  
 shepherdspurse  
 sowthistle, annual


fiddleneck, coast  
 filaree, broadleaf\*\*  
 filaree, redstem\*\*  
 filaree, whitestem\*\*  
 groundsel, common\*  
 henbit  
 mallow, little (malva, cheeseweed)  
 minerslettuce  
 nettle, burning  
 pigweed, redroot\*  
 redmaids  
 shepherdspurse  
 sowthistle, annual

\*Highest rate may be required for acceptable postemergence control.

\*\* Willowood OxyFlo 2EC at the 6-pint rate will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

**MINT (SPEARMINT AND PEPPERMINT)**

Mint (Spearmint and Peppermint) Grown on Mineral Soils		
Weed Control	Rate (pt/acre)	Specific Use Directions



<b>Preemergence Postemergence</b>	4 – 6	<b>Oregon and Washington (East of Cascades), California, Montana, Idaho, Nevada, South Dakota and Utah:</b> Apply from December through March when mint is dormant. When used postemergence (to weeds), add an 80% active ingredient nonionic surfactant at the rate of one quart per 100 gallons of spray volume and apply before weeds exceed a height of 4 inches. Late winter applications will provide maximum activity on summer weeds, but summer grass control may be inconsistent. For best results, fall-plowed fields should be harrowed to provide a smooth surface for application. In furrow-irrigated fields, corrugating must be done prior to application. Corrugating or harrowing will result in disturbance of treated soil or movement of untreated soil into treated areas, resulting in poor weed control.
<b>Preemergence</b>	2 – 3	<b>Peppermint (Western Oregon Willamette Valley):</b> Apply Willowood OxyFlo 2EC from November through February to dormant peppermint only. Treatments in January or February generally provide better residual preemergence control of annual broadleaf weeds. Full season weed control should not be expected from this treatment.

**Precautions:**

- Application must be made prior to emergence of new spring growth or severe crop injury may result.
- **In the Willamette valley, do not apply Willowood OxyFlo 2EC to mint that has been plowed.**
- Apply Willowood OxyFlo 2EC only to healthy stands of spearmint and peppermint. Do not apply to spearmint or peppermint weakened by disease, drought, flooding, excessive fertilizer, soil salts, previously applied pesticides, nematodes, insects, or winter injury, as severe injury may result.

**Crop-Specific Restrictions:**

- Do not make more than one application of Willowood OxyFlo 2EC per season.

**Key Weeds Controlled:**

bedstraw, catchweed	oats, wild*
bluegrass, annual*	orach, red
flixweed	pepperweed, yellowflower
groundsel, common	pigweed, redroot
lambsquarters, common	ryegrass, Italian*
lettuce, prickly (china lettuce)	shepherdspurse
mustard, blue (purple mustard)	sowthistle, annual
mustard, tumble (Jim hill mustard)	tansymustard
nightshade, hairy	thistle, Russian

\*Control of annual grasses is best obtained when Willowood OxyFlo 2EC is applied prior to emergence. Postemergence control of winter annual grasses is generally unsatisfactory if applications are made after the 1 to 2-leaf stage.

**Mint (Spearmint and Peppermint) Grown on Muck Soils: For Use Only on Mint Grown in Indiana, Michigan, Montana, North Dakota, South Dakota, and Wisconsin.**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	4 – 6	Note: Use directions in this section apply only to spearmint and peppermint grown on muck soils (organic matter content of 20% or greater). When used postemergence (to weeds), add an 80% active ingredient nonionic surfactant at the rate of one quart per 100 gallons of spray volume and apply before weeds exceed a height of 4 inches.

**Precautions:**

- Application must be made prior to emergence of new spring growth or severe crop injury may result.
- To avoid excessive crop injury, do not apply within 4 days of planting (sprigging) spearmint or peppermint.
- Apply Willowood OxyFlo 2EC only to healthy spearmint or peppermint. Do not apply to spearmint or peppermint that has been weakened by disease, nematodes, soil insects, or winter injury, as severe injury may result.

**Crop-Specific Restrictions:**

- Do not make more than one application of Willowood OxyFlo 2EC per season.

**Key Weeds Controlled:**

knotweed, prostrate  
pigweed, redroot  
purslane, common

**NON-CROP USE**

**(Non-Food-Producing, Non-Cultivated Agricultural or Non-Agricultural Areas, such as Highway and Utility Rights-of-Way, Industrial Sites, Tank Farms, Storage Areas, Airports, Fencerows, and Farmsteads)**

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b>	5 – 8	<p><b>Preemergence:</b> Use higher rate in rate range for longer residual control.</p> <p><b>Postemergence:</b> Use the lower rate in the rate range for control of susceptible weeds in the early postemergence stage less than 4 inches tall. Use the higher rate for weeds up to 12 inches tall. Application to weeds beyond the 4-inch stage may result in partial control.</p>
<b>Postemergence</b>	2 – 8	

**Tank Mixing:** Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.

- **Preemergence:** For broader-spectrum residual preemergence weed control, Willowood OxyFlo 2EC may be applied in tank mix combination with diuron (Karmex) or simazine.
- **Postemergence:** For additional postemergence control of susceptible grass and broadleaf weeds, Willowood OxyFlo 2EC may be applied in tank mix combination with paraquat (Gramoxone) or glyphosate.

**Site-Specific Restrictions:**

- Do not feed or allow animals to graze on any areas treated with Willowood OxyFlo 2EC.
- Do not apply more than 8 pints per acre in a single application.

**Key Weeds Controlled:**

### Preemergence

burclover  
cheeseweed (malva)  
fiddleneck, coast  
filaree, broadleaf  
filaree, redstem  
groundsel, common  
henbit  
knotweed, prostrate  
lambquarters, common  
lettuce, prickly  
pigweed, redroot  
purslane, common  
redmaids  
rocket, London  
shepherdspurse  
sowthistle, annual

### Postemergence

cheeseweed (malva)  
fiddleneck, coast  
filaree, broadleaf  
groundsel, common  
henbit  
minerslettuce  
nettle, burning  
pigweed, redroot  
purslane, common  
redmaids  
shepherdspurse  
sowthistle, annual

## ONIONS

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

**Cultural Considerations:** For maximum preemergence activity, the soil surface should be smooth and free of excessive trash (clippings, plant residues, etc.). Following application, cultural practices that result in redistribution or disturbance

of the soil surface or move untreated soil into treated areas will reduce weed control. For best results, make applications to established beds that are left undisturbed during the time period for which weed control is desired.

<b>Direct Seeded Onions: Postemergence Application</b>		
<b>Weed Control</b>	<b>Rate (per acre)</b>	<b>Specific Use Directions</b>
<b>Postemergence</b>	2 – 4 fl oz	<b>Northeastern States Including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont:</b> Apply Willowood OxyFlo 2EC at 2 to 4 fl oz per acre to seeded onions that have at least 3 true leaves using ground equipment. Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
<b>Postemergence</b>	0.5 – 1 pt	<b>Western States Including Arizona, California, Colorado, Idaho, Nevada, New Mexico, Oregon, Texas, Utah and Washington:</b> Apply Willowood OxyFlo 2EC at 0.5 to 1 pt per acre to seeded onions that have at least 2 true leaves using ground equipment. Multiple treatments at 0.5 to 1 pt per acre may be applied up to a maximum of 2 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
<b>Postemergence</b>	0.5 pt	<b>All Other States:</b> Apply Willowood OxyFlo 2EC at 0.5 pt per acre to seeded onions that have at least 2 true leaves using ground equipment. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.
<b>Postemergence</b>	(see above)	<b>Sprinkler Irrigation- All Except Northeastern States (Center Pivot, Portable Lateral or Solid Set):</b> Apply Willowood OxyFlo 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Follow the application directions and precautions for “Sprinkler Chemigation” given in the Chemigation section of this label.

<b>Transplanted Onions: Application Immediately before Planting</b>		
<b>Weed Control</b>	<b>Rate (per/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	1 – 2 pt	<b>Pre-transplant Application (Not for Use in Northeastern States or Western States):</b> Willowood OxyFlo 2EC may be applied as a broadcast or band application after completion of tillage operations, but before transplanting of onion plants. Transplanting should be accomplished with a minimum of soil disturbance and, for optimum weed control, soil surfaces should be left undisturbed after transplanting for the period for which weed control is desired. However, timely cultivation after weed emergence will assist in weed control. If less than 2 pt per acre was applied as a pre-transplant application, postemergence applications may be made as instructed for seeded onions. Do not exceed the maximum use rate of 2 pts per acre per use season as a result of multiple applications.

<b>Transplanted Onions: Application Immediately after Planting</b>		
<b>Application Timing for Target Weeds</b>	<b>Rate (per/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence</b>	up to 2 pts	<b>All States Except Northeastern States:</b> Transplanted onions are most tolerant of a postemergence application immediately after transplanting. An application of up to 2 pints per acre may be made within two days after transplanting. If less than 2 pints per acre is applied, a second application can be made two weeks or more after transplanting. Do not exceed the maximum use rate of 2 pints per acre of Willowood OxyFlo 2EC per season as a result of multiple applications.
<b>Preemergence</b>	2 – 4 oz	<b>Northeastern States including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont:</b> Multiple treatments at 2 to 4 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season.

**Onions- Use Precautions (Applicable to All Areas and Methods of Application):**

- Willowood OxyFlo 2EC can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the specified onion growth stage of the onion plants as specified in Specific Use Directions.
- Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects, nematodes or diseases.

**Onions-Crop-Specific Restrictions (Applicable to All Areas and Methods of Application):**

- In all states except Northeastern states, do not apply until direct seeded onion plants have at least two (2) fully developed true leaves. In the Northeastern states, do not apply until direct seeded onion plants have at least three (3) fully developed leaves. Application made prior to the specified growth stage may result in serious crop injury and is not recommended.
- Do not apply more than a total of 2 pints per acre of Willowood OxyFlo 2EC per use season as a result of multiple applications.
- Do not apply within 45 days of harvest.
- Do not apply Willowood OxyFlo 2EC as a preemergence treatment to direct seeded onions.
- Use only on dry bulb onions.
- Do not apply to onions grown for seed, except as instructed in separate use directions.
- For use in onions, do not mix Willowood OxyFlo 2EC with oils, surfactants, liquid fertilizers or pesticides except as specified on approved Willowood OxyFlo 2EC Supplemental Labeling.



**Key Weeds Controlled:****Postemergence**

canarygrass (annual)  
 eveningprimrose, cutleaf<sup>(a)</sup>  
 groundsel, common  
 mallow, little (malva)  
 nightshade, black  
 pigweed, prostrate<sup>(b)</sup>  
 pigweed, redroot<sup>(a,b)</sup>  
 puncturevine  
 purslane, common<sup>(a,b)</sup>  
 rocket, London  
 sage, lanceleaf  
 shepherspurse<sup>(b)</sup>  
 sowthistle, annual

(a) Weeds controlled when applied as a pre-transplant application. In addition, Willowood OxyFlo 2EC at the rate of 1 to 2 pints per acre will provide control/suppression of carpetweed, Pennsylvania smartweed, galinsoga, common lambsquarters, and wild mustard. Applications of Willowood OxyFlo 2EC to muck soils may result in partial control or suppression of the weeds listed.

(b) Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

**ONIONS GROWN FOR SEED**

**Agricultural Use Requirements:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Weed Control	Rate (per/acre)	Specific Use Directions
<b>Preemergence</b>	2 fl oz	<b>Northeastern States including Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island and Vermont:</b> Multiple treatments at 2 fl oz per acre may be applied up to a maximum of 2 pints (32 fl oz) per acre per use season. Prior to initial treatment, seeded onions must have <i>at least four (4) true leaves</i> . Multiple treatments at the aforementioned rate may be applied.

<b>Preemergence</b>	Up to 0.5 pt	<p><b>All other States:</b> Apply Willowood OxyFlo 2EC at up to 0.5 pt per acre to seeded onions that have <i>at least three (3) true leaves</i>. Multiple treatments at 0.5 pt per acre may be applied up to a maximum of 2 pints per acre per use season. For optimum postemergence control, apply when susceptible weeds are in the 2 to 4-leaf stage and actively growing.</p> <p><b>Sprinkler Irrigation- Portable Lateral or Solid Set:</b> Apply Willowood OxyFlo 2EC at the recommended broadcast application rate using sufficient irrigation to wet soil to a depth of 2 inches. Follow the application directions and precautions for “Sprinkler Chemigation” given in the Chemigation section of this label.</p>
---------------------	--------------	--

**Use Precautions:**

- **Notice:** Some varieties or inbred lines of onions may be more susceptible to Willowood OxyFlo 2EC. Care should be taken to ensure that the particular onion variety or line being grown is tolerant to Willowood OxyFlo 2EC. It is suggested that all onion varieties or lines be tested in limited areas to ensure an adequate level of crop tolerance prior to an application for postemergence weed control.
- Willowood OxyFlo 2EC can cause necrotic lesions, twisting, pigtailling or stunting of the onion plants. Injury will be more severe if applications are made immediately following or during cool, wet weather and/or if applications are made prior to the specified onion growth stage of the onion plants as specified in Specific Use Directions.
- Do not apply to onion plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, injury from previously applied pesticides, or injury due to insects or diseases.

**Crop-Specific Restrictions:**

- In all states, do not apply Willowood OxyFlo 2EC until the onions have reached the minimum leaf stage specified. Application prior to the recommended stage of development may result in serious injury and is not recommended.
- Do not apply more than a total of 2 pints per acre of Willowood OxyFlo 2EC during one use season.
- Do not apply within 60 days of harvest.
- For seeded onions, do not apply Willowood OxyFlo 2EC with oils, surfactants, liquid fertilizers or other pesticides except as specified in approved Willowood OxyFlo 2EC Supplemental Labeling.

**Key Weeds Controlled:**

**Postemergence**

canarygrass (annual)  
 eveningprimrose, cutleaf  
 groundsel, common  
 mallow, little (malva)  
 nightshade, black  
 pigweed, prostrate\*  
 pigweed, redroot\*  
 puncturevine  
 purslane, common\*  
 rocket, London  
 sage, lanceleaf  
 shepherdspurse  
 sowthistle, annual

\* Specific weeds controlled at rates recommended for use in northeastern states (see DOSAGE section).

## **PAPAYA**

**(For Use Only in Hawaii)**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	4	The initial application should occur no sooner than 4 months after transplanting or 6 months after direct seeding, and after the papaya has reached a minimum height of 4 feet. Applications may be repeated at approximate 4-month intervals.  Apply preemergence or postemergence to weeds. Increase the spray volume to assure adequate coverage of dense growth of emerged weeds. Willowood OxyFlo 2EC must be applied as a directed spray to the orchard floor beneath the papaya plants. Accurate, uniform placement of Willowood OxyFlo 2EC is essential for effective weed control and to minimize crop injury. Willowood OxyFlo 2EC must be applied using rigid precision ground sprayer equipment.  Postemergence applications may be made up to the 4 leaf stage of weed growth.
<b>Precautions:</b>		
<ul style="list-style-type: none"> <li>Do not allow the herbicide solution, spray, drift or mist to contact green bark, stems, fruit or foliage as injury may result.</li> <li>Do not use Willowood OxyFlo 2EC on papaya plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.</li> </ul>		
<b>Crop-Specific Restrictions:</b>		
<ul style="list-style-type: none"> <li>Do not apply more than 4 pints of Willowood OxyFlo 2EC per broadcast acre in a single directed spray or more than 12 pints per broadcast acre per year as a result of multiple applications.</li> <li>Do not apply Willowood OxyFlo 2EC within 1 day of harvest.</li> </ul>		

### **Key Weeds Controlled:**

amaranth, spiny  
purslane, common

spurge, garden

## **ROSES (FIELD-GROWN) IN CALIFORNIA**

### **Product Information**

Use this product as a post-directed application for control of certain broadleaf weeds in established rose plantings. Use Willowood OxyFlo 2EC for stool block ("permanent") plantings or rose plantings (field-grown, patio and tree roses) after bud grafted canes are at least 18-inches in length.

### **Crop Tolerance**

Apply this product only to roses with canes that are 18 inches or longer. Applications to rose plants with canes less than 18 inches in length can result in severe crop injury. Spray contact with foliage can cause severe crop injury and must be avoided. Leaves that are accidentally sprayed will exhibit necrotic spotting and may drop from the plant. Heavy splashing rain/irrigation or excessive soil moisture after application also can produce leaf cupping, crinkling, stunting or defoliation.

**Important:** When applied as directed, field-grown roses are tolerant to this product. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of roses or under all possible growing conditions. Exercise reasonable judgment with this product. Until familiar with results under user growing conditions, limit application of this product to a few plants in a small treated area to determine plant tolerance and extent of injury if such occurs, prior to initiating large-scale applications.

### **Method of Application**

Apply product in a minimum of 25 to 40 gallons of water per broadcast acre. Use a low-pressure sprayer with nozzles directed at the base of the rose plants. The use of spray shields also to avoid spray contact with rose foliage. To minimize spray drift, use the lowest spray pressure suitable for the application equipment being used.

This product is phytotoxic to plant foliage. Avoid drift to all other crops and nontarget areas. Do not apply when weather

conditions favor drift.

### Dosage

For postemergence weed control, apply Willowood OxyFlo 2EC at 2 to 4 pints (0.5 to 1.0 lb. active) per broadcast acre. Use the lower rate for the control of susceptible seedling weeds in the early postemergence stage up to the 4-leaf stage. Use the higher rate (1.0 lb. active) for weeds up to the 4-leaf stage. The addition of 1 quart of a nonionic surfactant per 100 gallons of spray mix will assist in spray coverage and wetting of weeds for postemergence coverage. Applications to weeds beyond the 4-leaf stage can result in partial control.

For preemergence weed control, apply this product at 2 to 4 pints (0.5 to 1.0 lb. active) per broadcast acre. Preemergence control is most effective when spray is applied to clean, weed-free soil surfaces. Seedling weeds are controlled as they come in contact with soil-applied herbicide during emergence.

### Weeds Controlled Preemergence

This product will provide control of the following weeds when applied at 2 to 4 pints (0.5 to 1.0 lb. active) per broadcast acre between the onset of germinating and the 4-leaf seedling growth stage. Partial control will likely result if weeds are beyond the 4-leaf seedling growth stage.

Beggarsticks, Nodding	Nightshade, Black
Cheeseweed (Malva)	Nightshade, Hairy
Morningglory, Annual	Pigweed, Redroot
Morningglory, Ivyleaf	Spurge, Spotted

### Weeds Controlled Postemergence

This product will provide control of the following weeds when applied at 2 to 4 pints (0.5 to 1.0 lb. active) per broadcast acre between the onset of germinating and the 4-leaf seedling growth stage. Partial control will likely result if weeds are beyond the 4-leaf seedling growth.

Cheeseweed (Malva)	Nightshade, Black
Morningglory, Annual	Nightshade, Hairy
Morningglory, Ivyleaf	Pigweed, Redroot

Willowood OxyFlo 2EC will also control or suppress many other annual broadleaf weeds.

### Field-Grown Roses – California Only

#### Specific Use Restrictions

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Do not apply more than 4 pints (1.0 lb. active) of this product per broadcast acre per application or more than 8 pints (2.0 lbs. active) per broadcast acre per growing season.
- Do not apply this product in an enclosed greenhouse structure, as injury to plant foliage may result. Tank mixtures of this product with oils, liquid fertilizers or other pesticides can result in enhanced crop response/injury and are the

responsibility of the user.

- Do not feed or graze animals on any area treated with Willowood OxyFlo 2EC.
- This product is phytotoxic to plant foliage. Do not apply when weather conditions favor drift to non-target areas.
- Do not apply this product to rose plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.
- Chemigation: Do not apply this product through any type of irrigation system.

**SOYBEANS**

**(Not for Use in California)**

<b>Soybeans – Early Preplant Application in Conservation Tillage Systems</b>		
<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence</b>	1.5-3	<p><b>Early Preplant Application:</b> Surface apply Willowood OxyFlo 2EC to the stale seedbed approximately 14 days before planting conservation tillage soybeans for postemergence and preemergence residual broadleaf control. Use a spray volume of 20 or more gallons per acre and increase the spray volume if growth of existing weeds is dense. Willowood OxyFlo 2EC at 2 to 3 pints provides early season suppression of annual grasses, but should not be relied upon as a basic grass herbicide. A planned program utilizing herbicides registered for early preplant, preemergence or postemergence grass control in soybeans is recommended.</p> <p>Use of ridge or slot planter or a similar planting implement that causes minimal soil disturbance is recommended. Movement or redistribution of surface soil will reduce herbicidal effectiveness.</p>

<b>Soybeans: No-Till (Double-Crop)</b>		
<b>Application Timing for Target Weeds</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	0.5-2	<p><b>Preemergence Application to Soybeans:</b> Applied preemergence, Willowood OxyFlo 2EC provides postemergence and residual preemergence control of susceptible broadleaf weeds. <b>Apply Willowood OxyFlo 2EC within one day after planting. Later applications may result in severe crop injury and are not recommended.</b> Apply in a minimum spray volume of 20 gallons per acre and increase spray volume if growth of existing weeds is dense.</p>

**Tank Mixing:** For enhanced postemergence control of existing grass and broadleaf weeds, Willowood OxyFlo 2EC may be tank mixed with paraquat (Gramoxone) or glyphosate. For extended residual control of annual grasses in no-till soybeans, Willowood OxyFlo 2EC may also be tank mixed with a residual grass herbicide such as Bronco Herbicide, Dual Magnum Herbicide, or Lasso Herbicide.

<b>Postemergence</b>	1	<b>Postemergence Directed Application:</b> Willowood OxyFlo 2EC may be applied as a post-directed application. Optimum control is achieved when Willowood OxyFlo 2EC is applied to seedling weeds not exceeding 4 true leaves (not counting cotyledon leaves) and actively growing. Use of an 80% nonionic surfactant cleared for application to growing crops at the rate of 2 pints per 100 gallons of spray is recommended whenever postemergence weed control is desired. <b>For postemergence application, soybeans must be a minimum 8 inches tall.</b> Use a minimum of 2 flat fan nozzles per row. Use branch lifters or shields to prevent excessive spray contact to the soybean plants. Do not use hollow cone nozzles.
----------------------	---	--

**Soybeans: Grown Under Conventional Tillage Systems**

<b>Application Timing for Target Weeds</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence Postemergence</b>	1-1.5	<b>Preemergence Application to Soybeans:</b> Willowood OxyFlo 2EC provides preemergence control of susceptible broadleaf weeds. <b>Apply Willowood OxyFlo 2EC within one day after planting. Later applications may result in severe crop injury and are not recommended.</b> Apply in a minimum spray volume of 20 gallons per acre and increase spray volume if growth of existing weeds is dense. The 1.5 pint per acre rate will assist in early season annual grass control but should not be relied upon as a basic grass herbicide. Willowood OxyFlo 2EC may also be applied as a preemergence application following a preplant incorporated grass herbicide treatment.

**Preemergence Tank Mixes (To Control Additional Grass and Broadleaf Weeds): Apply preemergence tank mixes of Willowood OxyFlo 2EC within one day after planting. Later applications may result in severe crop injury and are not recommended.**

- Willowood OxyFlo 2EC at 0.6 to 1.5 pints per acre may be applied preemergence to soybeans in tank mix with Dual Magnum Herbicide or Lasso Herbicide. Willowood OxyFlo 2EC may be applied alone as a preemergence application following a preplant incorporated grass herbicide application or as a tank mix in a preemergence application with Dual Magnum, or Lasso herbicides. Refer to the label of tank mix product for additional weeds controlled.
- Willowood OxyFlo 2EC at 0.6 to 0.8 pints per acre may be applied preemergence to soybeans in tank mix with 1 to 1.67 pints of Command 6EC herbicide. Refer to the label for Command 6EC for additional weeds controlled.

<b>Postemergence</b>	1	<b>Postemergence Directed Sprays:</b> Willowood OxyFlo 2EC may be applied as a post-directed application at 1 pint per acre. Optimum control is achieved when weeds do not exceed 4 true leaves and are actively growing (do not count cotyledon leaves). Use of an 80% nonionic surfactant cleared for application to growing crops at the rate of 2 pints per 100 gallons of spray is recommended whenever postemergence weed control is desired. <b>For postemergence application, soybeans must be a minimum of 8 inches tall.</b> Use a minimum of 2 flat fan nozzles per row. Use branch lifters or shields to prevent excessive spray contact to the soybean plants. Do not use hollow cone nozzles.
----------------------	---	---

**Postemergence Tank Mixes:** For broader spectrum control of broadleaf weeds, Willowood OxyFlo 2EC may be applied in tank mix with Butoxone Herbicide or Butyrac 200 Herbicide. Use 1 pint of Willowood OxyFlo 2EC with 1 pint of Butoxone or 0.7 to 0.9 pint of Butyrac 200 per acre. Refer to label of tank mix product for additional weeds controlled.

**Soybeans- Precautions (All Methods and Timings to Soybeans):**

- Soybeans are tolerant to preemergence and post-directed applications of Willowood OxyFlo 2EC at recommended rates; however, under certain conditions injury may occur. Heavy splashing rain shortly after crop emergence or cold, wet soil conditions during early growth stages can cause leaf cupping and crinkling. When injury occurs, it is generally limited to the first few leaves that develop after crop emergence. Soybeans recover from this injury and yields are not adversely affected. Soybeans accidentally sprayed during a post-directed application will exhibit necrotic spotting and injury to the soybean plant. Exercise care to avoid spray contact with the soybean leaves.

**Soybeans- Crop-Specific Restrictions:**

- **Tank Mixing:** Read and observe all label directions before using. Follow applicable use directions, precautions and limitations on the labels of the respective tank mix products. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive limitations must apply.
- Do not make more than two applications of Willowood OxyFlo 2EC per growing season.
- Do not apply more than 2 pints (0.5 lbs active) of Willowood OxyFlo 2EC per acre during one growing season as a result of preemergence application in no-till (double-crop) or conventional till soybeans, or post-directed in conventional till soybeans. If early preplant application is made, do not apply more than 3 pints (0.75 lb active) of Willowood OxyFlo 2EC per acre during one growing season.
- Do not apply a post-directed application of Willowood OxyFlo 2EC to soybeans after the initial appearance of blooms.

**Key Weeds Controlled (Willowood OxyFlo 2EC Alone):**

**Preemergence**

**Postemergence**

groundcherry, cutleaf\*  
 jimsonweed  
 lambsquarters, common  
 nightshade, American black\*  
 nightshade, black\*  
 pigweed, redroot  
 poinsettia, wild  
 shepherdspurse  
 sida, prickly (teaweed)  
 smartweed, Pennsylvania  
 sowthistle, common\*  
 velvetleaf

cocklebur, common  
 croton, tropic  
 groundcherry, cutleaf  
 groundcherry, Wright  
 jimsonweed  
 lambsquarters, common  
 morningglory, annual (up to 6 leaf)  
 mustard, wild  
 nightshade, American black  
 nightshade, black  
 nightshade, hairy  
 pigweed, redroot  
 poinsettia, wild\*  
 purslane, common  
 sesbania, hemp  
 shepherdspurse  
 sicklepod\*\*  
 sida, prickly (teaweed)\*  
 smartweed, Pennsylvania  
 velvetleaf

\* Multiple applications may be required for acceptable control.

\*\*Post-direct applications of Willowood OxyFlo 2EC will kill or suppress seedlings not exceeding the one true leaf stage.

## TARO

### (For Use Only in Hawaii)

For use only to dryland taro grown in Hawaii. Dryland taro is defined as taro grown without irrigation, or by using irrigation practices that do not result in run-off, irrigation return flow, or other loss of irrigation water from the production area. If irrigation is used, the water applied shall not exceed the field capacity of the soil.

Weed Control	Rate (pt/acre)	Specific Use Directions
Preemergence	2	<b>Preemergence to Taro and Weeds:</b> A single application of Willowood OxyFlo 2EC at the rate of 2 pints per acre may be applied within 1 week after transplanting but prior to emergence of taro plants.
Postemergence	1	<b>Postemergence to Taro and Weeds:</b> Willowood OxyFlo 2EC may be applied as a post-directed or band application at the rate of 1 pint per acre. Effective control of succulent weed seedlings in the 2-to 3-leaf stage can usually be obtained. Applications to weeds beyond the 3-leaf stage may result in partial control.

#### Precautions:

- Accurate, uniform placement of Willowood OxyFlo 2EC is essential for effective weed control and to minimize crop injury. Taro foliage receiving accidental spray or drift will be injured. Willowood OxyFlo 2EC must be applied using rigid precision ground sprayer equipment.
- Occasionally, after the use of Willowood OxyFlo 2EC, spotting, crinkling or flecking may appear on the leaves of the taro. Leaves that receive direct or indirect (drift) spray contact will be injured.
- Do not use Willowood OxyFlo 2EC on taro plantings that are weak, or under stress due to temperature, disease, fertilizer, nematodes, insects, pesticides, drought or excessive moisture.

#### Crop-Specific Restrictions:

- Do not apply more than 2 pints of Willowood OxyFlo 2EC per broadcast acre as a single preemergence application.
- Do not apply more than 1 pint of Willowood OxyFlo 2EC per acre in a single post-direct spray or more than 2 pints per acre per season as a result of multiple post-directed applications.
- Do not apply more than 4 pints of Willowood OxyFlo 2EC per acre per season as a result of preemergence and post-direct applications.
- Do not apply Willowood OxyFlo 2EC within 6 months of harvest of taro (corms, leaves).



**Key Weeds Controlled:**

amaranth, spiny  
 purslane, common  
 spurge, garden

**TREE FRUIT/NUT/VINE CROPS (Dormant Application)**

Almond, Apple, Apricot, Avocado, Beechnut, Brazil Nut, Bitternut, Cashew, Cherry, Chestnut, Chinquapin, Crab Apple, Date, Feijoa, Fig, Filbert, Grapes, Hickory Nut, Kiwi, Loquat, Macadamia Nut, Mayhaws, Nectarine, Olives, Peach, Pear, Pecan, Persimmon, Pistachio, Plum, Pomegranates, Prune, Quince, and Walnut

Weed Control	Rate (pt/acre)	Specific Use Directions
<b>Preemergence</b> (broadcast application)	5 – 6	Apply Willowood OxyFlo 2EC in a minimum of 20 gallons of water per acre. Use higher spray volumes to ensure thorough coverage in high densities of emerged weeds or heavy trash. Sprays should be directed to the soil and the base of dormant trees or vines.
(banded application)	5 – 8	<b>In California</b> , Willowood OxyFlo 2EC may be applied as an over-the-top or directed spray to dormant nonbearing grape plantings. The use of a low-pressure sprayer is suggested. Do not apply over-the-top to grape plantings that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, injury from previously applied pesticides, or injury due to insects, nematodes, or diseases, as severe crop injury may result.
<b>Postemergence</b> (broadcast application)	2 – 6	Apply in a spray volume of 40 or more gallons per acre. For optimum control, apply when weeds are at seedling stage of growth.
(banded application)	2 – 8	The lower rate in the rate range (2 pints per acre) is recommended for the control of susceptible seedling weeds in the early postemergence stage up to the 4-leaf stage. Higher rates (up to 6 pints per acre) may be used for weeds up to the 6-leaf stage. Applications to weeds beyond the 6-leaf stage may result in partial control.

**Tank Mixing:** Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply. See labels of tank mix partners to determine suitability and use rates for various crops.

- **Postemergence:** for broader spectrum postemergence control of listed grass and broadleaf weeds, Willowood OxyFlo 2EC may be applied in tank mix with paraquat (Gramoxone) or glyphosate. These herbicides may also be added to preemergence tank mixes for enhanced control of existing weeds.
- **Preemergence:** For broad-spectrum preemergence control of susceptible grass and broadleaf weeds in listed tree fruit, nut or vine plantings, Willowood OxyFlo 2EC may be applied in tank mix with napropamide (Devrinol herbicide), diuron (Karmex herbicide), pronamide (Kerb® herbicide), simazine, norfluazon (Solicam herbicide) or oryzalin (Surflan herbicide).

**Chemigation (All States):** For dormant season application using sprinkler (low-volume (micro sprinkler)), drip (trickle), and flood (basin) irrigation systems, apply Willowood OxyFlo 2EC at the specified rate per acre. Follow applicable directions in the Chemigation section of this label when making applications using irrigation systems.

**Precautions:**

- Willowood OxyFlo 2EC or any of the combinations recommended on this label should be applied to only healthy growing trees or vines.
- Avoid direct plant contact. Direct spray toward the base of tree or vines unless specific use recommendations allow over-the-top application.

**Crop-Specific Restrictions:**

- In all states, unless otherwise specified, do not apply Willowood OxyFlo 2EC during the period between bud swell and completion of final harvest or when fruit/nuts are present. Willowood OxyFlo 2EC may be applied upon completion of final harvest.
- **In Arizona and California, Willowood OxyFlo 2EC may be applied during the period following completion of final harvest up to February 15 (February 1st in the Coachella Valley, California). Applications made after these calendar dates, but prior to bud swell, may result in significant crop injury and are the responsibility of the user.**
- For banded applications, up to 8 pints per acre of Willowood OxyFlo 2EC per use season may be applied within the treated band. Do not apply more than a maximum of 6 pints per acre per use season on a broadcast basis.
- Do not apply to grapes or kiwi established less than 3 years unless vines are on a trellis wire a minimum of 3 feet above the soil surface.
- Do not apply to grapes or kiwi that are not staked or trellised unless vines are free standing.

**Key Weeds Controlled (Arizona and California):****Preemergence**

burclover  
 cheeseweed (malva)  
 fiddleneck, coast  
 filaree, broadleaf  
 filaree, redstem  
 filaree, whitestem  
 groundsel, common  
 henbit  
 knotweed, prostrate  
 lambsquarters, common  
 lettuce, prickly  
 pigweed, redroot  
 purslane, common  
 redmaids  
 rocket, London  
 shepherdspurse  
 sowthistle, annual

**Postemergence**

cheeseweed (malva)  
 fiddleneck, coast  
 filaree, broadleaf\*  
 filaree, redstem\*  
 filaree, whitestem\*  
 groundsel, common  
 henbit  
 minerslettuce  
 nettle, burning  
 pigweed, redroot  
 redmaids  
 shepherdspurse  
 sowthistle, annual

\* Willowood OxyFlo 2EC at the 6-pint rate will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control.

**Key Weeds Controlled (All Other States Except Arizona and California)****Preemergence****Postemergence**

camphorweed  
 cudweed, narrowleaf  
 eveningprimrose, cutleaf\*  
 groundcherry, cutleaf  
 jimsonweed  
 lambsquarters, common  
 nightshade, American black  
 nightshade, black  
 pepperweed, Virginia  
 pigweed, redroot  
 poinsettia, wild  
 sida, prickly  
 smartweed, Pennsylvania  
 sowthistle, annual  
 spurge, prostrate  
 spurge, spotted  
 velvetleaf

balsamapple  
 cocklebur, common  
 cudweed, narrowleaf\*\*  
 eveningprimrose, cutleaf\*\*\*  
 groundcherry, cutleaf  
 groundcherry, Wright  
 jimsonweed  
 lambsquarters, common  
 morningglory, annual  
 nightshade, American black  
 nightshade, black  
 pepperweed, Virginia  
 pigweed, redroot  
 poinsettia, common  
 sesbania, hemp  
 shepherdspurse  
 sida, prickly (teaweed)  
 smartweed, Pennsylvania  
 sowthistle, annual  
 velvetleaf

\*Highest rate and/or multiple applications may be required for acceptable control.

\*\*Maximum 0.5-inch diameter

\*\*\*Highest rate and/or multiple applications may be required for acceptable control.

**GRAPES (Non-Dormant Application)**

**(California Only)**

Willowood OxyFlo 2EC may be applied as a directed spray or, for supplemental preemergence weed control, through low-volume sprinkler (micro sprinkler) or drip irrigation systems for control or suppression of listed broadleaf weeds in non-dormant grapes (raisin and wine grapes only). Willowood OxyFlo 2EC may also be applied to all grapes (raisin, table, and wine) as a dormant season application. Refer to Tree fruit/Nut/Vine Crops (Dormant Application) section above for use directions for dormant season application to grapes.

Weed Control	Rate (pt/acre)	Specific Use Directions
--------------	----------------	-------------------------

<b>Preemergence</b>	2	Willowood OxyFlo 2EC may be applied preemergence or postemergence to weeds either as a directed spray in a minimum spray volume of 20 gallons per acre or through low-volume sprinkler (micro sprinkler) or drip irrigation systems. Repeat applications may be required. Applications may be made from completion of bloom up to 14 days before harvest.  When applied as a postemergence directed spray, add 1 quart 80% active nonionic surfactant cleared for application to growing crops per 100 gallons of spray. Sprays should be directed to the soil and the base of vines.
<b>Postemergence</b>	1 – 2	

**Tank Mixing:**

- When applied as a directed postemergence spray using ground equipment, Willowood OxyFlo 2EC may be applied in tank mix with paraquat (Gramoxone) or glyphosate in a minimum spray volume of 10 gallons per acre. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.

**Chemigation:** Follow chemigation instructions in Product Information section.

- **Low Volume Sprinkler (Microsprinkler) and Drip (Trickle) Irrigation:** Apply only through low-volume sprinkler or drip systems designed to uniformly distribute irrigation water beneath the canopy. Meter Willowood OxyFlo 2EC at a continuous rate during the middle 1/3 of the irrigation period and discontinue application during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. Use of Willowood OxyFlo 2EC through low-volume sprinklers or drip emitters helps to reduce the “ring effect” of weed escapes in areas around sprinklers or emitters where previously applied broadcast or directed treatments begin to break down.

**Precautions:**

- **Crop Tolerance:** The use of Willowood OxyFlo 2EC may result in varying degrees of injury to non-dormant grapes. Grape foliage will typically exhibit injury symptoms from direct or indirect (spray drift, soil contact) exposure. This injury may result in necrosis, reddening, cupping or crinkling of grape leaves. The grape plant will continue to grow normally. Grape leaves that are immature or expanding at the time of contact with Willowood OxyFlo 2EC are the most susceptible to foliage injury. Grapes may exhibit some small blemishes (spots or flicks) on the fruit.
- Willowood OxyFlo 2EC is phytotoxic to plant foliage. Avoid drift to all other crops and nontarget areas. Do not apply when weather conditions favor drift.

**Crop-Specific Use Restrictions:**

- The total amount of Willowood OxyFlo 2EC applied during one season (from completion of final harvest through dormancy to non-dormant use covered by this section) cannot exceed 6 pints per acre as a result of multiple applications in any given area (broadcast, banded, or within the wetted area of the low-volume sprinkler or drip irrigation system.)
- Do not apply within 14 days of harvest.
- Do not initiate application of Willowood OxyFlo 2EC in non-dormant grapes until the completion of the bloom period.
- Do not apply to grapes established less than 3 years unless vines are either on a trellis wire a minimum of 3 feet above the soil surface, or protected by grow tubes.
- Willowood OxyFlo 2EC should be applied only by ground application equipment or through low-volume sprinkler (micro sprinkler) or drip (trickle) irrigation systems.
- Apply Willowood OxyFlo 2EC as a non-dormant application to wine grapes or raisin grapes only.

**Key Weeds Controlled or Suppressed:**

**Preemergence**

**Postemergence**

burclover  
cheeseweed, malva  
fiddleneck, coast  
groundsel, common  
henbit  
knotweed, prostrate  
lambsquarters, common  
minerslettuce  
mustard, black  
nettle, burning  
nightshade, black  
pigweed, redroot  
purslane, common  
redmaids  
rocket, London  
sowthistle, annual

cheeseweed, (malva)  
fiddleneck, coast  
groundsel, common  
henbit  
minerslettuce  
morningglory species, annual  
mustard, black  
nettle, burning  
nightshade, black  
pigweed, redroot  
purslane, common  
redmaids  
rocket, London  
sowthistle, annual

## **GRAPES-NONBEARING**

### **(WASHINGTON ONLY)**

#### **Dormant-Nonbearing Grapes In Washington**

##### **Product Information**

This product is effective as a preemergence and/or postemergence herbicide for the control of certain annual grasses and broadleaf weeds in dormant nonbearing grape plantings. The most effective postemergence weed control is achieved when Willowood OxyFlo 2EC is applied to seedling weeds.

##### **Crop Tolerance**

When this product is applied according to labeled directions, crop response (leaf cupping, crinkling and necrosis) and

stunting can be observed on new emerging growth. Vines typically outgrow this condition and develop normally. Some varieties or root stocks of grapes may be more susceptible to this product. Ensure that the particular grape variety being grown is tolerant to Willowood OxyFlo 2EC.

### Timing

Applications of this product must not be made after budswell. In the fall, Willowood OxyFlo 2EC can be applied after the grape plantings are dormant or frost has occurred. Applications prior to dormancy or frost occurring can result in significant crop injury and are the responsibility of the user. Apply this product to dormant vines, prior to budswell. The closer the grapes are to bud-break at application, the greater the crop response can be to Willowood OxyFlo 2EC. Do not apply this product after the buds start to swell.

### Method of Application

Apply product in a minimum of 40 gallons of water per acre. Use higher volumes to ensure adequate coverage in high densities of emerged weeds or heavy trash. Best preemergence results are achieved when spray is applied to a relatively weed-free established berm or soil surface. Apply this product as an over-the-top or directed spray to dormant nonbearing grape plantings. Use a low-pressure sprayer.

### Dosage

Apply Willowood OxyFlo 2EC for postemergence control at 2 to 8 pints (0.5 to 2.0 lbs. active) per broadcast acre. For preemergence susceptible weeds, use 5 to 8 pints (1.25 to 2.0 lbs. active) per broadcast acre.

### Weeds Controlled Preemergence

Apply 5 to 8 pints (1.25 to 2.0 lbs. active) of this product per broadcast acre.

Burclover	Lambsquarters, Common
Cheeseweed (Malva)	Lettuce, Prickly
Fiddleneck, Coast	Pigweed, Redroot
Filaree, Broadleaf	Purslane, Common
Filaree, Redstem	Redmaids
Flixweed	Rocket, London
Groundsel, Common	Shepherdspurse
Henbit	Sowthistle, Annual
Knotweed, Prostrate	Spurge, Prostrate
	Velvetleaf

### Weeds Controlled Postemergence

Apply 2 to 8 pints (0.5 to 2.0 lbs. active) of this product per broadcast acre to weeds up to 4 inches high. Applications to weeds beyond this 4-inch stage may result in partial control.

Cheeseweed (Malva)	Miner's Lettuce
Cocklebur, Common	Mustard, Wild
Fiddleneck, Coast	Nettle, Burning
Filaree, Redstem*	Nightshade, Black
Flixweed	Nightshade, Hairy
Groundsel, Common	Pigweed, Redroot
Henbit	Shepherdspurse
Ladysthumb	Sowthistle, Annual

\*Willowood OxyFlo 2EC at the 8-pint rate (2.0 lbs. active) will provide control of filaree not exceeding the 4-inch stage. Applications to filaree beyond the 4-inch stage may result in partial control. When postemergence weed control is desired, add an 80% active nonionic surfactant cleared for application to growing crops at the rate of 2 pints per each 100 gallons of spray.

### Nonbearing Grapes – Washington Only

#### Dormant Only

#### Specific Use Restrictions

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- DO NOT APPLY TO NONDORMANT GRAPES. Read and observe all label directions before using.
- Do not apply more than 8 pints (2.0 lbs. active) per broadcast acre of this product in one season.
- This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife, and fish. Willowood OxyFlo 2EC must not be used under this label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult other sections of this label for additional restrictions and precautions to protect aquatic organisms.
- Tank mixtures of this product with other pesticides may result in enhanced crop response/injury and are the responsibility of the user.
- Do not apply to grape plantings that are under stress due to drought, flooding, excessive fertilizer or soil salts, storage conditions, wind injury, hail, injury from previously applied pesticides, or injury due to insects, nematodes, or diseased, as severe crop injury may result.
- Chemigation: Do not apply this product through any type of irrigation system.
- A 25-ft. vegetative buffer strip must be maintained between all areas treated with this product and lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial ponds.

### SUCKER CONTROL IN NON-DORMANT GRAPES

#### (Washington and Oregon Only)

#### (Grapes for Wine and Processing Only)

Application Timing for Sucker Control	Rate (pt/acre)	Specific Use Directions
Grape suckers less than 12 inches in length.	1 – 2	Apply Willowood OxyFlo 2EC in a three-foot band directed towards newly emerging suckers at the base of the grapevine. The highest rate and/or a second application may be required to achieve an acceptable level of control/suppression of grape suckers. Avoid spray contact on flowers, grape clusters, or fruit. Use mounted nozzles to deliver the spray solution. Thorough spray coverage of sucker growth is essential for optimal activity. Use a spray volume of 50 or more gallons per acre (broadcast basis).

**Tank Mixing:** For enhanced postemergence sucker activity, a tank mixture of Willowood OxyFlo 2EC with either glufosinate (Rely Herbicide) or paraquat (Gramoxone) can be used. Apply at the recommended rates and growth stages in a manner described on the respective labels. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.

#### Precautions:

- The use of Willowood OxyFlo 2EC may result in varying degrees of injury to non-dormant grapes. Grape foliage will typically exhibit injury symptoms from direct or indirect (spray drift or soil contact) exposure. This injury may result in necrosis, reddening, cupping or crinkling of grape leaves. The grape plant will continue to grow normally. Leaves that are immature or expanding at the time of contact with Willowood OxyFlo 2EC are the most susceptible to injury. Grape fruit may exhibit some small blemishes (spots or flecks) on the fruit.

**Crop-Specific Restrictions:**

- The total amount of Willowood OxyFlo 2EC applied during one crop year (dormant and non-dormant) cannot exceed 6 pints per acre as a result of multiple applications in any given area (broadcast or banded).
- Willowood OxyFlo 2EC should be applied only by ground application equipment.
- Apply Willowood OxyFlo 2EC as a non-dormant application for sucker control only to wine or processed grapes.
- Do not apply Willowood OxyFlo 2EC within 60 days of harvest.

**APRICOTS, NECTARINES, OLIVES, PEACHES, PLUMS AND PRUNES****(CALIFORNIA ONLY)****Nondormant Application to Apricots, Nectarines, Olives, Peaches, Plums and Prunes in California****Product Information**

This product provides effective postemergence control of cheeseweed (Malva), Fleabane, and Marestalk (Horseweed), young broadleaf weed seedlings in non-dormant apricots, nectarines, olives, peaches, plums and prunes. For enhanced postemergence activity against these target weeds as well as other weed species, tank mix this product with either paraquat (Parazone 3SL) or glyphosate (Glyphogan) to increase the spectrum of weed control by either of these tank mix partners. Compatibility of each mixture must be established before tank mixing and application must be applied by ground equipment. Follow all precautions and restrictions on the labeling of the products to be tank mixed.

**Dosage**

Use this product for postemergence suppression of the target weeds at 0.5 to 1 pint (0.125 to 0.25 lb. active) per broadcast acre when applied to susceptible weed seedlings less than 4 inches in height. Repeat applications may be required. Do not exceed 6 pints (1.5 lbs. active) of this product during the non-dormant stage of apricots, nectarines, olives, peaches, plums and prunes. For a broader spectrum of grass weeds and broadleaf weeds control in the tree row middles, a tank mixture of Willowood OxyFlo 2EC with either paraquat (Parazone 3SL) or glyphosate (Glyphogan) can be used. Read and follow the labeling of either the paraquat (Parazone 3SL) or glyphosate (Glyphogan) pesticide product which is to be tank mixed with this product.

**Method of Application**

**Ground Application:** Apply a minimum spray volume of 10 gallons of water per acre. Use higher volumes to ensure adequate coverage in high densities of emerged weeds or heavy trash. Use conventional low-pressure ground spray equipment with flat fan spray nozzles at 20 to 40 psi. Position an off-center nozzle at the end of the boom. Calibrate spray equipment carefully before each use.

**Chemigation Application:** Apply this product only through flood (basin) irrigation systems, or low-volume sprinkler (microsprinkler) and drip (trickle) irrigation systems designed to distribute irrigation water beneath the tree canopy. For additional information on these systems, see the **Chemigation Instructions** section of this label.

**Cultural Considerations for All Applications:** In order to provide maximum effectiveness of preemergence activity of this product, the berm or soil surface must be level, smooth, and free of crop or weed trash (decaying leaves, clippings, dead weeds, etc.). Remove leaves and trash by blowing the area to be treated or by thoroughly mixing the trash into the soil through cultivation prior to herbicide applications.

Cultural practices that result in redistribution or disturbance of the soil surface after treatment will decrease the herbicidal effectiveness of this product. Cutting water furrows or cultivations that mix untreated soil into treated areas will also reduce the effectiveness of the treatment. For best results, apply to established berms or soil surfaces that are left undisturbed during the time period for which weed control is desired.

**Apricots, Nectarines, Olives, Peaches, Plums and Prunes in California****Nondormant Application****Specific Use Restrictions**

In addition to the following, also observe the use restrictions listed at the beginning of this label.

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturer labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- When applied as a non-dormant treatment, this product can only be applied to apricots, peaches, nectarines, plums and prunes after May 1. Willowood OxyFlo 2EC can only be applied as a non-dormant treatment to olives after bloom.
- Do not apply this product within 14 days of harvest of fruit.
- Do not apply more than 6 pints (1.5 lbs. a.i.) per broadcast acre of this product during the non-dormant season



- Apply this product only to healthy trees.
- Direct spray toward the base of the tree. Avoid direct herbicide contact with foliage and fruit.

### **PISTACHIOS, WALNUTS, ALMONDS (CALIFORNIA AND ARIZONA ONLY)**

#### **(Non-Dormant Application)**

<b>Weed Control</b>	<b>Rate (pt/acre)</b>	<b>Specific Use Directions</b>
<b>Preemergence</b>	5 – 6	<b>Preemergence:</b> For residual weed control of listed weeds.
<b>Postemergence</b>	1 – 2	<b>Postemergence (Suppression):</b> Apply to seedling weeds less than 4 inches in height. Repeat applications may be required.
	2 – 6	<b>Postemergence (Cleanup):</b> Contact (postemergence) control for cleanup sprays and preharvest applications. Apply to seedling weeds less than 4 inches in height. Applications to weed seedlings beyond the 4-inch stage may result in partial control.

**CALIFORNIA ONLY:** Willowood OxyFlo 2EC may be applied at a rate of no more than 5 pts/acre (1.25 lbs. a.i.) before February 15, and no more than 0.5 pt/acre (0.125 lb. a.i.) up to 30 days before harvest and/or no more than 0.5 pt/acre (0.125 lb. a.i.) between 30 and 15 days before harvest. Do not apply more than 6 pints of Willowood OxyFlo 2EC or 1.5 lb. oxyfluorfen a.i. per broadcast acre during the non-dormant season.

**Tank Mixing:** For broader spectrum grass and broadleaf weed control in tree row middles, Willowood OxyFlo 2EC may be tank mixed with either paraquat (Gramoxone) or glyphosate. Refer to Mixing Directions section for Tank Mixing Precautions. Follow applicable use directions, precautions, and limitations on the respective product labels. In interpreting the labels of tank mixed products, the most restrictive label limitations must apply.

**Chemigation:** Follow chemigation instructions in Product Information section.

**Flood (Basin) Irrigation:** For flood (basin) irrigation systems, meter continuously into the water during the entire irrigation period. Best weed control results are obtained when a uniform distribution and flow of irrigation water is maintained over level land. Irrigation water treated with Willowood OxyFlo 2EC must be contained on the treated area until the water is absorbed by the soil.

**Low Volume Sprinkler (Microsprinkler) and Drip (Trickle) Irrigation:** Apply only through low-volume sprinkler or drip systems designed to uniformly distribute irrigation water beneath the tree canopy. Applications should be made prior to weed emergence; otherwise postemergence activity may be inconsistent due to uneven coverage. Meter Willowood OxyFlo 2EC at a continuous rate during the middle 1/3 of the irrigation period and discontinue application during the final 1/3 of the irrigation period to ensure proper flushing of the irrigation system. Use of Willowood OxyFlo 2EC through low-volume sprinklers or drip emitters helps to reduce the “ring effect” of weed escapes in areas around sprinklers or emitters where previously applied broadcast or directed treatments begin to break down.

#### **Precautions:**

- Direct spray toward the base of trees. Avoid direct contact with foliage or nuts.
- Willowood OxyFlo 2EC should be applied only to healthy growing trees.

#### **Crop-Specific Use Restrictions:**

- When applied as a non-dormant treatment, Willowood OxyFlo 2EC can only be applied to pistachio plantings between May and 7 days prior to harvest.
- When applied as a non-dormant treatment, Willowood OxyFlo 2EC can only be applied to almond plantings between April 1 and September 30 and to walnut plantings between May 1 and September 30.
- Do not apply Willowood OxyFlo 2EC within 7 days of harvest of pistachios.
- Do not apply Willowood OxyFlo 2EC within 30 days of harvest of almonds (AZ only) or within 15 days of harvest of almonds (CA only).
- Do not apply Willowood OxyFlo 2EC within 7 days of harvest of walnuts.
- Do not apply more than 6 pints of Willowood OxyFlo 2EC or 1.5 lb. oxyfluorfen active ingredient per acre during the non-dormant season.

#### **Key Weeds Suppressed and/or Controlled**

cheeseweed, (malva)  
 fiddleneck, coast  
 filaree, broadleaf  
 filaree, redstem  
 filaree, whitestem  
 groundsel, common  
 henbit  
 minerslettuce

morningglory species, annual  
 mustard, black  
 nettle, burning  
 pigweed, redroot  
 purslane, common  
 redmaids  
 rocket, London  
 sowthistle, annual

**Additional Weeds Controlled in Tank Mix with Glyphosate or Paraquat**

barnyardgrass  
 bluegrass, annual  
 chickweed, common

horseweed (marestail)  
 rocket, London  
 ryegrass, Italian

**WINDBREAKS AND SHELTERBELTS**

(For Use Only in Minnesota, North Dakota, South Dakota and Wyoming)

Weed Control	Rate (pt/acre)	Specific Use Directions
<p><b>Preemergence</b> <b>Postemergence</b></p>	<p>4 – 6</p>	<p>Willowood OxyFlo 2EC may be applied as a broadcast, banded or post-directed spray. Preemergence control is most effective when spray is applied to clean, weed-free soil surfaces. Pre-transplant applications must be made after completion of soil preparation but prior to transplanting. Transplanting should be completed with minimal soil disturbance. For optimum weed control results, treated soil surfaces should be left undisturbed during the time period for which weed control is desired.</p> <p><b>Postemergence Weed Control:</b> For best results, apply before 4-leaf stage for broadleaf weeds or 2-leaf stage for grass weeds.</p> <p><b>Conifers:</b> Willowood OxyFlo 2EC can be applied pre-transplant, post-directed or postemergence (over-the-top) to conifers. Postemergence or post-directed applications should be applied prior to budbreak or after new growth foliage has hardened off and new terminal buds have formed.</p> <p><b>Deciduous Hardwoods:</b> Willowood OxyFlo 2EC has exhibited selectivity to many deciduous species when applied pre-transplant or as a post-directed spray prior to budbreak.</p>

**Precautions:**

- **Important:** Some varieties or cultivars of conifers or deciduous species listed may be susceptible to Willowood OxyFlo 2EC. Care should be taken to ensure that the particular variety to be sprayed with Willowood OxyFlo 2EC is tolerant. For unfamiliar species, it is suggested that Willowood OxyFlo 2EC be tested on a limited number of plants prior to large-scale application.
- Occasionally after the use of Willowood OxyFlo 2EC, a spotting, crinkling or flecking may appear on the leaves of the deciduous species. Leaves that receive direct or indirect (drift) spray contact will be injured. Deciduous species typically rapidly outgrow these symptoms and develop normally.
- Application after budbreak may result in injury to deciduous species and is not recommended. If non-dormant application is required, apply only after foliage has fully expanded and hardened off. Avoid direct or indirect spray contact with the foliage by applying to the soil surface as a directed spray.
- Apply Willowood OxyFlo 2EC only to healthy deciduous and/or conifer trees. Do not apply Willowood OxyFlo 2EC to conifers or deciduous trees that have been weakened or under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result.

**Specific Use Restrictions for Shelterbelts:**

- Do not apply more than 6 pints of Willowood OxyFlo 2EC per acre in a single application or more than 18 pints per acre per year.

**Key Broadleaf Weeds Controlled:**

buckwheat, wild  
 burclover  
 carpetweed  
 dock, curly  
 groundcherry, cutleaf  
 groundcherry, Wright  
 groundsel, common  
 henbit  
 jimsonweed  
 knotweed, prostrate  
 kochia  
 ladythumb  
 lambsquarters, common  
 lettuce, prickly  
 mallow, little  
 mayweed  
 mustard, blue  
 mustard, tumble

mustard, wild  
 nettle, burning  
 nightshade, black  
 nightshade, hairy  
 oats, wild  
 orach, red  
 pepperweed, yellow flower  
 pigweed, prostrate  
 pigweed, redroot  
 purslane, common  
 rocket, London  
 shepherdspurse\*  
 smartweed, Pennsylvania  
 sowthistle, annual  
 tansymustard  
 thistle, Russian (seedling)  
 velvetleaf

\* The highest rate or multiple applications may be required for acceptable control.

**Key Grasses Controlled:**

barnyardgrass	foxtail, giant
bluegrass, annual	goosegrass
crabgrass, large	witchgrass

**Willowood OxyFlo 2EC may be applied to numerous conifer and deciduous species, including the following:**

**Conifer Species**

<b>Common Name</b>	<b>Scientific Name</b>
douglas-fir	<i>Pseudotsuga menziesii</i>
fir	
grand	<i>Abies grandis</i>
fraser	<i>Abies fraseri</i>
noble	<i>Abies procera</i>
hemlock	
eastern hemlock	<i>Tsuga Canadensis</i>
western hemlock	<i>Tsuga heterophylla</i>
pine	
Austrian	<i>Pinus nigra</i>
eastern white	<i>Pinus strobes</i>
jack	<i>Pinus banksiana</i>
Himalayan	<i>Pinus graffithii</i>
loblolly	<i>Pinus taeda</i>
lodgepole	<i>Pinus contorta</i>
longleaf	<i>Pinus palustris</i>
monterey	<i>Pinus radiate</i>
mugo	<i>Pinus mugo</i>
ponderosa	<i>Pinus ponderosa</i>
scotch	<i>Pinus sylvestris</i>
shortleaf	<i>Pinus echinata</i>
slash	<i>Pinus elliotti</i>
Virginia	<i>Pinus virginiana</i>

spruce	
blue	<i>Picea pungens</i>
dwarf Alberta	<i>Picea glauca conica</i>
Norway	<i>Picea abies</i>
Sitka	<i>Picea sitchensis</i>
Arborvitae	<i>Thuja occidentalis</i> <i>Thuja orientalis</i>
juniper	<i>Juniperus chinensis</i> <i>Juniperus horizontalis</i> <i>Juniperus procumbens</i> <i>Juniperus sabina</i> <i>Juniperus scopulorum</i>
red cedar	<i>Juniperus virginiana</i>
yew	<i>Taxus</i> spp.

### Deciduous Hardwood Species

Common Name	Scientific Name
ash	<i>Fraxinus</i> spp.
crabapple	<i>Malus</i> spp.
eucalyptus	<i>Eucalyptus</i> spp.
lilac	<i>Syringa vulgaris</i>
maple, black	<i>Acer nigrum</i>
oak, northern red	<i>Quercus rubra</i>
olive, Russian	<i>Elaeagnus angustifolia</i>
poplar (cottonwood)	<i>Populus</i> spp.
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product above 40°F (5°C).

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Willowood, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Willowood, LLC and Seller harmless for any claims relating to such factors.

Willowood, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Willowood, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Willowood, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WILLOWOOD, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WILLOWOOD, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Willowood, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Willowood, LLC.

EPA 20111108