

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR WILLOWOOD ETHOFUMESATE 4SC AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

This labeling must be in the possession of the user at the time of pesticide application.

“Label” as used in this supplemental labeling refers to the label booklet for Willowood Ethofumesate 4SC and this supplement.

This supplemental labeling expires on July 30, 2020 and must not be used or distributed after this date.

Willowood ETHOFUMESATE 4SC

FOR USE IN SUGAR BEETS

EPA Reg. No. 87290-1

WILLOWOOD ETHOFUMESATE 4SC + GLYPHOSATE TANK MIX FOR WEED CONTROL IN SUGAR BEETS

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRODUCT INFORMATION

Willowood Ethofumesate 4SC is a selective herbicide for use in sugar beets for the control of the weed species listed on the product label. It provides effective control of these weeds for up to 10 weeks following application.

Residual control of weeds is dependent upon soil moisture conditions; rate of herbicide used, and soil texture. The activity of Willowood Ethofumesate 4SC in the soil is reduced as the soil texture becomes finer and organic matter increases.

POSTEMERGENCE APPLICATION

WILLOWOOD ETHOFUMESATE 4SC + GLYPHOSATE (TANK MIX)

Postemergence application of Willowood Ethofumesate 4SC plus glyphosate can improve control of weeds on the label including glyphosate-resistant biotypes.

POSTEMERGENCE APPLICATIONS

Apply Willowood Ethofumesate 4SC + glyphosate to Roundup Ready sugar beets only.

Apply Willowood Ethofumesate 4SC + glyphosate to sugar beets having greater than 2 true leaves.

Apply Willowood Ethofumesate 4SC + glyphosate to small weeds.

Apply Willowood Ethofumesate 4SC+ glyphosate in a single or multiple application(s).

The greater the number of Willowood Ethofumesate 4SC + glyphosate applications, the greater the weed control.

Include the following spray adjuvants to tank mixes of Willowood Ethofumesate 4SC + glyphosate: ammonium sulfate plus methylated seed oil (MSO) at 1.25 to 2 pints/A or high surfactant methylated seed oil (HSMOC) at 1.0 to 2 pints/A or nonionic surfactant (NIS) at 0.25% v/v. Do not include a crop oil concentrate adjuvant with this mixture.

WILLOWOOD USA 

Willowood, LLC
385 Interlocken Crescent, Suite #240
Broomfield, CO 80021

DOSAGE TABLE 5

WILLOWOOD ETHOFUMESATE 4SC + GLYPHOSATE DOSAGE FOR POSTEMERGENCE APPLICATIONS

Sugar Beet Stage	WILLOWOOD ETHOFUMESATE 4SC PINTS/ACRE BROADCAST
> 2 true leaves to 45-days prior to harvest	1/2 - 8

Willowood Ethofumesate 4SC + Glyphosate Postemergence Use

Precautions

Ethofumesate 4SC + glyphosate applied at greater than 5 pints/A to fields having coarse-textured low organic matter soils will likely cause crop response.

Sugar beet injury is possible when applying Willowood Ethofumesate 4SC + glyphosate at greater than 5 pints/A to small sugar beets.

Removing MSO, HSMOC, and NIS from tank mixes with metolachlor, dimethenamid, acetochlor, triflusulfuron-methyl and clopyralid should decrease crop response, but weed control may be reduced.

Tank mixes of metolachlor, dimethenamid, acetochlor, triflusulfuron-methyl and/or clopyralid plus adjuvants with Willowood Ethofumesate 4SC + glyphosate will cause crop response and likely yield loss, especially with increasing rates of Willowood Ethofumesate 4SC.

Sugar beet injury can occur when Willowood Ethofumesate 4SC is applied preemergence and metolachlor, dimethenamid or acetochlor are mixed with Willowood Ethofumesate 4SC + glyphosate postemergence.

Sugar beet injury will be enhanced if Willowood Ethofumesate 4SC + glyphosate is applied following a preplant application of eptam.

Sugar beet injury can be enhanced if the crop is under stress.

Restrictions

Do not apply greater than a total of 8 pints/A (4 lb a.i./A) of Willowood Ethofumesate 4SC in a single or multiple postemergence application(s) per year. Do not apply greater than 8 pints/A (4 lb a.i./A) for the entire year for all applications. Do not apply more than 2 postemergent applications at a rate of 2 lb a.i./A.

Aerial Application Rate Restriction: Do not apply more than a total of 1.5 lbs. a.i. (3 pints of product) per acre per application with aircraft.

Allow at least 10 days between postemergence applications.

Willowood Ethofumesate 4SC may be applied up to 45 days before harvest. For all products mixed with Willowood Ethofumesate 4SC, follow the most restrictive pre-harvest interval listed on the product label.

DOSAGE TABLE

WILLOWOOD ETHOFUMESATE 4SC + GLYPHOSATE - MAXIMUM DOSAGE AND NUMBER OF APPLICATIONS FOR POSTEMERGENCE APPLICATIONS

	Annual Rate			Annual Application Rate	
	Max. Single Rate of Ethofumesate, lb. AI/A	Max. # Applications of Willowood Ethofumesate 4SC per Year	Max. Ethofumesate Rate, lb. AI/A	Max. No. crop cycles per year	Max. Rate, Ethofumesate lb. AI/A/year)
4.0	4	4.0	1	4.0	

EPA 20171207



Willowood, LLC
385 Interlocken Crescent, Suite #240
Broomfield, CO 80021